arXiv:0708.0656v1 [math.ST] 5 Aug 2007

A MULTIVARIATE CENTRAL LIMIT THEOREM FOR RANDOMIZED ORTHOGONAL ARRAY SAMPLING DESIGNS IN COMPUTER EXPERIMENTS[1]

By Wei-Liem Loh

National University of Singapore

Let $f : [0,1)^d \to \mathbb{R}$ be an integrable function. An objective of many computer experiments is to estimate $\int_{[0,1)^d} f(x)dx$ by evaluating f at a finite number of points in $[0,1)^d$. There is a design issue in the choice of these points and a popular choice is via the use of randomized orthogonal arrays. This article proves a multivariate central limit theorem for a class of randomized orthogonal array sampling designs [Owen (1992a)] as well as for a class of OA-based Latin hypercubes [Tang (1993)].

1 Introduction

Let X be a random vector uniformly distributed on the d-dimensional unit hypercube $[0,1)^d$ and f be an integrable function from $[0,1)^d$ to $\mathbb{R}$. An objective of many computer experiments [see, for example, McKay, Conover and Beckman (1979), Stein (1987), Sacks, Welch, Mitchell and Wynn (1989) and Santner, Williams and Notz (2003)] is to estimate

$$\mu = Ef(X) = \int_{[0,1)^d} f(x)dx, \tag{1}$$

using a finite number of function evaluations. It is well known that as the dimension d increases, Monte Carlo methods and (deterministic) equidistribution methods become competitive and ultimately dominant. Indeed Davis and Rabinowitz (1984), Chapter 5.10, consider $d > 15$ to be a high enough dimensionality that sampling or equidistribution methods are indicated.

For definiteness, let n, d, q and t be positive integers such that $t \leq d$. An orthogonal array of strength t is a matrix of n rows and d columns with elements taken from the set of symbols $\{0, 1, \ldots, q-1\}$ such that in any $n \times t$ submatrix, each of the q^t possible rows occurs the same number of times. The class of all such arrays is denoted by OA(n, d, q, t).

---

[1] *AMS* 2000 *subject classifications.* Primary 62E20; secondary 60F05, 65C05.

Key words and phrases. Computer experiment, multivariate central limit theorem, numerical integration, OA-based Latin hypercube, randomized orthogonal array, Stein's method.

Comprehensive accounts of orthogonal arrays can be found in the books by Raghavarao (1971) and Hedayat, Sloane and Stufken (1999).

Owen (1992a), (1994) and Tang (1993) independently proposed the use of randomized orthogonal arrays in computer experiment sampling designs. The main attraction of these designs is that they, in contrast to simple random sampling, stratify on all t-variate margins simultaneously. A class of randomized orthogonal array sampling designs proposed by Owen (1992a) is as follows. Let

(a) $A \in \mathrm{OA}(q^t, d, q, t)$ where $a_{i,j}$ denotes the (i,j)th element of A,

(b) $\pi_1, \ldots, \pi_d$ be random permutations of $\{0, \ldots, q-1\}$, each uniformly distributed on all the $q!$ possible permutations,

(c) $\{U_{i,j} : i = 1, \ldots, q^t, j = 1, \ldots, d\}$, be $[0,1)$ uniform random variables,

(d) and all the $U_{i,j}$'s and π_k's are independent.

We randomize the symbols of A by applying the permutation π_j to the jth column of A, $j = 1, \ldots, d$. This gives us another orthogonal array A^* such that its (i,j)th element satisfies $a^*_{i,j} = \pi_j(a_{i,j})$. An orthogonal array based sample of size q^t (taken from $[0,1)^d$) is defined to be $\{X_1, \ldots, X_{q^t}\}$ where for $i = 1, \ldots, q^t$, $X_i = (X_{i,1}, \ldots, X_{i,d})'$,

$$X_{i,j} = \frac{a^*_{i,j} + U_{i,j}}{q}, \qquad \forall j = 1, \ldots, d. \tag{2}$$

For $t \geq 2$, Tang (1993) observed that the above sampling designs may not stratify well on s-variate margins if $s < t$. He suggested modified designs that stratify on t-variate margins as well as 1-variate margins simultaneously. He called these designs OA-based Latin hypercubes. Finally, Owen (1997a), (1997b), in a series of articles, proposed the use of scrambled nets. Given $t \in \mathbb{Z}^+$, the scrambled nets stratify on s-variate margins whenever t/s is a positive integer.

A class of OA-based Latin hypercubes can be constructed as follows. Let $A \in \mathrm{OA}(q^t, d, q, t)$. As before, we randomize its symbols to obtain the orthogonal array A^*. Then for each column of A^*, we replace the q^{t-1} positions with entry k by a random permutation (with each such permutation having an equal probability of being chosen) of $\{kq^{t-1}, kq^{t-1} + 1, \ldots, (k+1)q^{t-1} - 1\}$, for all $k = 0, \ldots, q-1$. After the replacement is done for all d columns of A^*, the newly obtained matrix, say A^{**}, satisfies $A^{**} \in \mathrm{OA}(q^t, d, q^t, 1)$.

One version of OA-based Latin hypercubes that was considered by Owen (1997a), page 1906, is of the form $\{Y_1, \ldots, Y_{q^t}\}$ where for $i = 1, \ldots, q^t$, $Y_i = (Y_{i,1}, \ldots, Y_{i,d})'$,

$$Y_{i,j} = \frac{a^{**}_{i,j} + U_{i,j}}{q^t}, \quad \forall j = 1, \ldots, d, \tag{3}$$

$\{U_{i,j} : i = 1, \ldots, q^t, j = 1, \ldots, d\}$ are $U[0,1)$ random variables independent of one another and all other permutations, and $a^{**}_{i,j}$ denotes the (i,j)th element of A^{**}. The class of OA-based Latin hypercubes proposed by Tang (1993) requires one more level of randomization where the columns of A^{**} are randomized. We denote the resulting matrix by A^{***}. Tang's OA-based Latin hypercubes can be expressed as $\{Y^*_1, \ldots, Y^*_{q^t}\}$ where for $i = 1, \ldots, q^t$, $Y^*_i = (Y^*_{i,1}, \ldots, Y^*_{i,d})'$,

$$Y^*_{i,j} = \frac{a^{***}_{i,j} + U_{i,j}}{q^t}, \quad \forall j = 1, \ldots, d, \tag{4}$$

$\{U_{i,j} : i = 1, \ldots, q^t, j = 1, \ldots, d\}$ are, as before, $U[0,1)$ random variables independent of one another and all other permutations, and $a^{***}_{i,j}$ denotes the (i,j)th element of A^{***}. We note that $\{Y_1, \ldots, Y_{q^t}\}$ and $\{Y^*_1, \ldots, Y^*_{q^t}\}$ are Latin hypercube samples [see, for example, McKay, Conover and Beckman (1979) and Owen (1992b)].

The estimators for μ in (1) that we are concerned with are

$$\hat{\mu}_{oas} = q^{-t} \textstyle\sum_{i=1}^{q^t} f(X_i),\ \hat{\mu}_{oal} = q^{-t} \sum_{i=1}^{q^t} f(Y_i), \text{ and } \hat{\mu}^*_{oal} = q^{-t} \sum_{i=1}^{q^t} f(Y^*_i), \tag{5}$$

where the X_i's, Y_i's and Y^*_i's are as in (2), (3) and (4) respectively. It is easily seen that $\hat{\mu}_{oas}$, $\hat{\mu}_{oal}$ and $\hat{\mu}^*_{oal}$ are all unbiased estimators for μ. For simplicity, we write $\sigma^2_{oas} = \mathrm{Var}(\hat{\mu}_{oas})$, $\sigma^2_{oal} = \mathrm{Var}(\hat{\mu}_{oal})$ and $\sigma^{*2}_{oal} = \mathrm{Var}(\hat{\mu}^*_{oal})$.

In this article, we shall assume that $t = 2$. This significantly simplifies the notation as well as the theoretical arguments that follow. Also as Owen (1992a) and Tang (1993) noted, orthogonal arrays of strength $t = 2$ lead to the most economical sample size q^2. This is important in practice especially when q is large. The following theorem is due to Owen (1992a) and Tang (1993).

Theorem 1. *Let $d \geq 3$, f be a bounded continuous function on $[0,1)^d$ and $\hat{\mu}_{oas}$, $\hat{\mu}^*_{oal}$ be as in (5) with $A \in OA(q^2, d, q, 2)$. Then as $q \to \infty$, we have*

$$q^2 \sigma^2_{oas} = \int_{[0,1)^d} f^2_{rem}(x) dx + o(1) \quad \text{and} \quad q^2 \sigma^{*2}_{oal} = \int_{[0,1)^d} f^2_{rem}(x) dx + o(1),$$

where for all $x = (x_1, \ldots, x_d)' \in [0,1)^d$, $1 \leq j \leq d$, $1 \leq k < l \leq d$,

$$f_j(x_j) = \int_{[0,1)^{d-1}} [f(x) - \mu] \prod_{1 \leq k \leq d : k \neq j} dx_k,$$

$$\begin{aligned} f_{k,l}(x_k, x_l) &= \int_{[0,1)^{d-2}} [f(x) - \mu - f_k(x_k) - f_l(x_l)] \prod_{1 \leq j \leq d: j \neq k,l} dx_j, \\ f_{rem}(x) &= f(x) - \mu - \sum_{j=1}^{d} f_j(x_j) - \sum_{1 \leq k < l \leq d} f_{k,l}(x_k, x_l). \end{aligned} \tag{6}$$

Theorem 1 implies that (i) the asymptotic variances of $\hat{\mu}_{oas}$ and $\hat{\mu}^*_{oal}$ are always less than or equal to the asymptotic variance of an analogous estimator based on a simple random sample of the same size, (ii) they are dramatically smaller if the integrand f can be approximated by a sum of bivariate functions, and (iii) $\sigma^2_{oas} \sim \sigma^{*2}_{oal}$ if $\int_{[0,1)^d} f^2_{rem}(x)dx > 0$. Tang (1993), page 1395, further showed that $\sigma^{*2}_{oal} \leq \sigma^2_{oas}$ if f is additive.

The aim of this article is to study the asymptotic distributions of $\hat{\mu}_{oas}$, $\hat{\mu}_{oal}$ and $\hat{\mu}^*_{oal}$. For instance, such a result will be useful in the construction of confidence intervals for μ.

Definition. A function $f : [0,1)^d \rightarrow \mathbb{R}$ is smooth with a Lipschitz continuous mixed partial of order d if there exist finite constants $B \geq 0$ and $\beta \in (0,1]$ such that

$$\sup_{j_1,\ldots,j_d \in \{0,\ldots,d\}: j_1+\cdots+j_d = d} \left| \frac{\partial^d}{\partial x_1^{j_1} \ldots \partial x_d^{j_d}} f(x) - \frac{\partial^d}{\partial x_1^{j_1} \ldots \partial x_d^{j_d}} f(y) \right| \leq B \|x - y\|^\beta,$$

$\forall x, y \in [0,1)^d$ where $\|.\|$ is the usual Euclidean norm. We shall now state the main result of this article, the proof of which is deferred to the Appendix.

Theorem 2. *Suppose $d \geq 3$ and $f : [0,1)^d \rightarrow \mathbb{R}$ is smooth with a Lipschitz continuous mixed partial of order d such that $\int_{\mathbb{R}^d} f^2_{rem}(x)dx > 0$. Define $W_{oas} = (\hat{\mu}_{oas} - \mu)/\sigma_{oas}$, $W_{oal} = (\hat{\mu}_{oal} - \mu)/\sigma_{oal}$ and $W^*_{oal} = (\hat{\mu}^*_{oal} - \mu)/\sigma^*_{oal}$ with $A \in OA(q^2, d, q, 2)$. Then W_{oas}, W_{oal} and W^*_{oal} each converges in law to the standard normal distribution as $q \rightarrow \infty$.*

The remainder of this article proceeds as follows. In Section 2 we shall first establish base q expansions for $\{X_1, \ldots, X_{q^2}\}$ and $\{Y_1, \ldots, Y_{q^2}\}$. The main point here is that the difference between these two base q expansions is of order $O(1/q)$. Following Owen (1997a), a d-dimensional base q Haar multiresolution analysis is applied to f and an ANOVA decomposition of f is obtained. This ANOVA decomposition facilitates much of the theoretical analysis that ensue.

In Section 3, a proxy statistic W for W_{oas} and W_{oal} is introduced. Proposition 2 shows that to prove the asymptotic normality of W_{oas} and W_{oal} as $q \rightarrow \infty$, it suffices to prove that W is asymptotically normal. Stein (1972) proposed a powerful and general method for obtaining a bound for the error in the normal approximation to the distribution of a sum of dependent random variables. Since then, Stein's method has found considerable applications in combinatorics, probability and statistics [see Stein (1986)]. We shall use the multivariate normal version of Stein's method as given in Götze (1991) and Bolthausen and Götze (1993). In particular, Theorem 3 establishes a multivariate central limit theorem for the "components" of W under the conditions of Theorem 2. This result is needed in the proof of the latter theorem. Finally, the Appendix contains the proof of Theorem 2 as well as some more technical results that used in this article.

We would like to add that Loh (1996) has established the asymptotic normality of $\hat{\mu}_{oas}$ when $d = 3$ and $t = 2$ under moment conditions on f. However the approach in Loh (1996), which uses directly the univariate version of Stein's method, does not seem to be extendable to $d \geq 4$. For example, the inequality (11) in Loh (1996) is valid for $d = 3$ but not for $d \geq 4$.

We conclude the Introduction with a note on notation. In this article, the indicator function is denoted by $\mathcal{I}\{.\}$ and if x is a vector, then x' is its transpose. $\|.\|$ denotes the Euclidean norm in $\mathbb{R}^p$ where p is either $d-2$ or d (depending on the context).

2 ANOVA decomposition

We shall first establish base q expansions for randomized orthogonal array samples as well as for OA-based Latin hypercubes. Let $A \in \mathrm{OA}(q^2, d, q, 2)$, $a_{i,j}$ be the (i,j)th element of A and

$$\{\pi_j, \pi_{j;b}, \pi_{i,j,k} : i = 1, \ldots, q^2, j = 1, \ldots, d, b = 0, \ldots, q-1, k = 2, 3, \ldots\} \tag{7}$$

be a set of mutually independent random permutations of $\{0, 1, \ldots, q-1\}$, where each of these permutations is uniformly distributed over its $q!$ possible values. We observe that the randomized orthogonal array sample $X_1, \ldots, X_{q^2}$ in (2) can be expressed as $X_i = (X_{i,1}, \ldots, X_{i,d})'$ where

$$X_{i,j} = \sum_{k=1}^{\infty} x_{i,j,k} q^{-k}, \qquad \forall i = 1, \ldots, q^2, j = 1, \ldots d, \tag{8}$$

$x_{i,j,1} = \pi_j(a_{i,j})$ and $x_{i,j,k} = \pi_{i,j,k}(0)$ for all $k \geq 2$. Let $A^{**} \in \mathrm{OA}(q^2, d, q^2, 1)$ be as in Section 1 with $t = 2$. Since $0 \leq a^{**}_{i,j}/q^2 < 1$, we observe that $a^{**}_{i,j}/q^2 = \sum_{k=1}^{\infty} b_{i,j,k} q^{-k}$ for suitable integers $0 \leq b_{i,j,1}, b_{i,j,2} \leq q-1$ and $b_{i,j,k} = 0$ for all $k \geq 3$. Owen (1997a), page 1907, observed that an OA-based Latin hypercube defined as in (3) has the form $\{Y_i = (Y_{i,1}, \ldots, Y_{i,d})' : i = 1, \ldots, q^2\}$, where

$$Y_{i,j} = \sum_{k=1}^{\infty} y_{i,j,k} q^{-k} \tag{9}$$

and for $1 \leq i \leq q^2$, $1 \leq j \leq d$, $y_{i,j,1} = \pi_j(a_{i,j})$, $y_{i,j,2} = \pi_{j;a_{i,j}}(b_{i,j,2})$, $y_{i,j,k} = \pi_{i,j,k}(0)$ for all $k \geq 3$. We observe from (8) and (9) that $\sup_{1 \leq i \leq q^2, 1 \leq j \leq d} |X_{i,j} - Y_{i,j}| \leq (q-1)/q^2$.

Let $f : [0,1)^d \to \mathbb{R}$ be a square integrable function. Inspired by Owen (1997a), we apply a d-dimensional base q Haar multiresolution analysis to f. More precisely, for any integer $k \geq 0$, let $\mathcal{Y}_k$ denote the linear span of the functions $\{\psi_{k,t,c} : t = 0, 1, \ldots,$ and $c = 0, \ldots, q-1\}$ where

$$\psi_{k,t,c}(x) = q^{(k+1)/2}\mathcal{I}\{x \in [\frac{qt+c}{q^{k+1}}, \frac{qt+c+1}{q^{k+1}})\} - q^{(k-1)/2}\mathcal{I}\{x \in [\frac{t}{q^k}, \frac{t+1}{q^k})\},$$

$\forall x \in [0,1)$. We observe that the functions in $\mathcal{Y}_k$ are constant on $[tq^{-k-1}, (t+1)q^{-k-1})$ and integrate to zero over $[tq^{-k}, (t+1)q^{-k})$. Next let $\mathcal{U}_0$ denote the space of functions that are constant on $[0,1)$ and

$$\mathcal{U}_k = \{g + g_0 + \cdots + g_{k-1} : g \in \mathcal{U}_0, g_j \in \mathcal{Y}_j, j = 0, \ldots, k-1\}, \qquad \forall k = 1, 2, \ldots.$$

Then it is well known that $\bigcup_{k=0}^{\infty}\mathcal{U}_k$ is dense in $L^2([0,1))$ and $\bigcap_{k=0}^{\infty}\mathcal{U}_k = \mathcal{U}_0$. We further observe from Owen (1997a), page 1897, that a typical basis function for $L^2([0,1)^d)$ is of the form $\prod_{r=1}^{l}\psi_{k_{j_r},t_{j_r},c_{j_r}}(x_{j_r})$ for all $(x_1, \ldots, x_d)' \in [0,1)^d$, where $1 \leq j_1 < \cdots < j_l \leq d$, and $k_{j_r} \geq 0$, $0 \leq t_{j_r} \leq q^{k_{j_r}} - 1$, $0 \leq c_{j_r} \leq q-1$ whenever $1 \leq r \leq l$. Here by convention, an empty product (that is $l = 0$) is taken to be 1. Hence for each $f \in L^2([0,1)^d)$, it follows from (6.6) of Owen (1997a), page 1898, that

$$\begin{aligned} f(x) &= \mu + \sum_{l=1}^{d} \sum_{1 \leq j_1 < \cdots < j_l \leq d} (\sum_{k_{j_1}=0}^{\infty} \sum_{t_{j_1}=0}^{q^{k_{j_1}}-1} \sum_{c_{j_1}=0}^{q-1}) \cdots (\sum_{k_{j_l}=0}^{\infty} \sum_{t_{j_l}=0}^{q^{k_{j_l}}-1} \sum_{c_{j_l}=0}^{q-1}) \\ &\qquad \times \langle f, \prod_{r=1}^{l}\psi_{k_{j_r},t_{j_r},c_{j_r}}\rangle \prod_{r=1}^{l}\psi_{k_{j_r},t_{j_r},c_{j_r}}(x_{j_r}), \qquad \text{a.e. } x \in [0,1)^d, \end{aligned} \tag{10}$$

where μ is as in (1) and

$$\langle f, \prod_{r=1}^{l}\psi_{k_{j_r},t_{j_r},c_{j_r}}\rangle = \int_{[0,1)^d} f(x)[\prod_{r=1}^{l}\psi_{k_{j_r},t_{j_r},c_{j_r}}(x_{j_r})]dx. \tag{11}$$

Without loss of generality, we can assume that equality in (10) holds for all $x \in [0,1)^d$ since changing the value of f on a set of Lebesgue measure zero will not alter the value of μ. For simplicity let

$$\{U[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d] : 0 \leq \tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} \leq q-1, u_j \geq 0, 1 \leq j \leq d\}$$

be a set of mutually independent random vectors where each $U[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d]$ has the uniform distribution on the d-dimensional interval $\prod_{j=1}^{d}[\sum_{k=1}^{u_j}\tilde{c}_{j,k}q^{-k}, q^{-u_j} + \sum_{k=1}^{u_j}\tilde{c}_{j,k}q^{-k})$. Here $\sum_{k=1}^{u_j}\tilde{c}_{j,k}q^{-k} = 0$ if $u_j = 0$. Furthermore we assume that the above U's are independent of π's [defined as in (7)]. For nonnegative integers $u_1^*, u_1, \ldots, u_d^*, u_d$, we write

(i) $(u_1^*, \ldots, u_d^*) \preceq (u_1, \ldots, u_d)$ if and only if $u_j^* \leq u_j$ for all $j = 1, \ldots, d$,

(ii) $(u_1^*, \ldots, u_d^*) \prec (u_1, \ldots, u_d)$ if and only if $u_j^* \leq u_j$ for all $j = 1, \ldots, d$ with at least one strict inequality.

The following construction establishes an ANOVA decomposition of $Ef \circ U = Ef(U)$ where E denotes expectation. For integers $u_j \geq 0$, $0 \leq \tilde{c}_{j,k} \leq q-1$, $1 \leq j \leq d, k \geq 1$, define recursively

$$\begin{aligned} & \nu_{u_1,\ldots,u_d}[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d] \\ = \; & Ef \circ U[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d] \\ & - \sum_{u_1^*,\ldots,u_d^* : (0,\ldots,0) \preceq (u_1^*,\ldots,u_d^*) \prec (u_1,\ldots,u_d)} \nu_{u_1^*,\ldots,u_d^*}[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j^*} : 1 \leq j \leq d], \end{aligned} \tag{12}$$

and hence

$$\begin{aligned} & \lim_{u_1,\ldots,u_d \to \infty} Ef \circ U[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d] \\ = \; & \sum_{u_1,\ldots,u_d \geq 0} \nu_{u_1,\ldots,u_d}[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d]. \end{aligned}$$

Writing $|u| = \sum_{j=1}^d \mathcal{I}\{u_j \geq 1\}, u = (u_1, \ldots, u_d)'$, such that $1 \leq j_1 < \cdots < j_{|u|} \leq d$ and $u_j \geq 1$ if and only if $j \in \{j_1, \ldots, j_{|u|}\}$, it follows from (10) that $\nu_{u_1,\ldots,u_d}$ can be written down explicitly as $\nu_{0,\ldots,0}[\quad] = \mu$ if $|u| = 0$ and

$$\begin{aligned} & \nu_{u_1,\ldots,u_d}[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d] \\ = \; & \Big(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1} \sum_{c_{j_1}=0}^{q-1}\Big) \cdots \Big(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1} \sum_{c_{j_{|u|}}=0}^{q-1}\Big) \langle f, \prod_{l=1}^{|u|} \psi_{u_{j_l}-1, t_{j_l}, c_{j_l}} \rangle \\ & \times E\{\prod_{l=1}^{|u|} \psi_{u_{j_l}-1, t_{j_l}, c_{j_l}} \circ U_{j_l}[\tilde{c}_{j,1}, \ldots, \tilde{c}_{j,u_j} : 1 \leq j \leq d]\} \\ = \; & \Big(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1} \sum_{c_{j_1}=0}^{q-1}\Big) \cdots \Big(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1} \sum_{c_{j_{|u|}}=0}^{q-1}\Big) \langle f, \prod_{l=1}^{|u|} \psi_{u_{j_l}-1, t_{j_l}, c_{j_l}} \rangle \\ & \times \prod_{l=1}^{|u|} \psi_{u_{j_l}-1, t_{j_l}, c_{j_l}} \Big(\sum_{k=1}^{u_{j_l}} \tilde{c}_{j_l,k} q^{-k}\Big), \end{aligned} \tag{13}$$

if $|u| \geq 1$. Here U_{j_l} denotes the j_lth co-ordinate of U and the last equality uses the fact that $\psi_{u_{j_l}-1, t_{j_l}, c_{j_l}}$ is constant on $[tq^{-u_{j_l}}, (t+1)q^{-u_{j_l}})$ for an arbitrary but fixed integer t. An important consequence of the ANOVA decomposition (that will be applied repeatedly in the sequel) is if $u_k \geq 1$ for some $1 \leq k \leq d$, then

$$\sum_{\tilde{c}_{k,u_k}=0}^{q-1} \nu_{u_1,\ldots,u_d}[\tilde{c}_{j,1},\ldots,\tilde{c}_{j,u_j} : 1 \leq j \leq d] = 0. \tag{14}$$

Writing $W_{oal} = (\hat{\mu}_{oal} - \mu)/\sigma_{oal}$, we observe from (5), (9), (12), (13) and (14) that

$$\begin{aligned} W_{oal} &= \frac{1}{q^2 \sigma_{oal}} \sum_{i=1}^{q^2} [f(Y_i) - \mu] \\ &= \frac{1}{q^2 \sigma_{oal}} \sum_{i=1}^{q^2} \sum_{u_1,\ldots,u_d : (0,\ldots,0) \prec (u_1,\ldots,u_d)} \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \pi_{j;a_{i,j}}(b_{i,j,2}), \\ &\qquad \pi_{i,j,3}(0),\ldots,\pi_{i,j,u_j}(0) : 1 \leq j \leq d] \\ &= \frac{1}{q^2 \sigma_{oal}} \sum_{i=1}^{q^2} \sum_{u_1,\ldots,u_d \geq 0 : u_1+\cdots+u_d \geq 3} \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \pi_{j;a_{i,j}}(b_{i,j,2}), \\ &\qquad \pi_{i,j,3}(0),\ldots,\pi_{i,j,u_j}(0) : 1 \leq j \leq d]. \end{aligned} \tag{15}$$

Writing $W_{oas} = (\hat{\mu}_{oas} - \mu)/\sigma_{oas}$ and in a similar manner to (15), we have

$$\begin{aligned} W_{oas} &= \frac{1}{q^2 \sigma_{oas}} \sum_{i=1}^{q^2} \sum_{u_1,\ldots,u_d \geq 0 : u_1+\cdots+u_d \geq 3} \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \\ &\qquad \pi_{i,j,2}(0),\ldots,\pi_{i,j,u_j}(0) : 1 \leq j \leq d] \\ &\quad + \frac{1}{q^2 \sigma_{oas}} \sum_{i=1}^{q^2} \sum_{1 \leq k \leq d : u_k = 2, u_l = 0 \forall l \neq k} \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \\ &\qquad \pi_{i,j,2}(0),\ldots,\pi_{i,j,u_j}(0) : 1 \leq j \leq d]. \end{aligned} \tag{16}$$

For brevity of notation, we write in the sequel

$$\begin{aligned} & \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \pi_{j;a_{i,j}}(b_{i,j,2}), \pi_{i,j,3}(0),\ldots,\pi_{i,j,u_j}(0) : 1 \leq j \leq d] \\ = \; & \nu^*_{u_{j_1},\ldots,u_{j_{|u|}}}[\pi_{j_1}(a_{i,j_1}), \pi_{j_1;a_{i,j_1}}(b_{i,j_1,2}), \pi_{i,j_1,3}(0),\ldots,\pi_{i,j_1,u_{j_1}}(0);\ldots; \\ & \qquad \pi_{j_{|u|}}(a_{i,j_{|u|}}), \pi_{j_{|u|};a_{i,j_{|u|}}}(b_{i,j_{|u|},2}), \pi_{i,j_{|u|},3}(0),\ldots,\pi_{i,j_{|u|},u_{j_{|u|}}}(0)], \end{aligned} \tag{17}$$

$\nu^*[.] = \nu^*_{u_{j_1},\ldots,u_{j_{|u|}}}[.]$ if $u_{j_1} = \cdots = u_{j_{|u|}} = 1$, and

$$\sigma^2 = E\Big\{\{\frac{1}{q^2} \sum_{i=1}^{q^2} \sum_{0 \leq u_1,\ldots,u_d \leq 1 : |u| \geq 3} \nu^*[\pi_{j_1}(a_{i,j_1});\ldots;\pi_{j_{|u|}}(a_{i,j_{|u|}})]\}^2\Big\}, \tag{18}$$

where $1 \leq j_1 < \cdots < j_{|u|} \leq d$ are exactly those coordinates of $u = (u_1,\ldots,u_d)'$ in which $u_j \geq 1$ and $|u|$ denotes the cardinality of that set. We end this section with the following proposition.

Proposition 1. *Let $f : [0,1)^d \to \mathbb{R}$ be smooth with a Lipschitz continuous mixed partial of order d. Then*

$$\begin{aligned}
\sigma^2_{oal} &= q^{-2} \sum_{0 \le u_1,\ldots,u_d \le 1 : |u| \ge 3} E\{\nu^*[\pi_{j_1}(a_{1,j_1}); \ldots; \pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} + O(q^{-3}), \\
\sigma^2_{oas} &= \sigma^2_{oal} + O(q^{-3}) \text{ and } \sigma^2 = \sigma^2_{oal} + O(q^{-3}) \text{ as } q \to \infty.
\end{aligned}$$

Proof. Since $E(W^2_{oal}) = 1$, we observe from (15) that

$$\begin{aligned}
\sigma^2_{oal} &= \frac{1}{q^4} \sum_{i_1=1}^{q^2} \sum_{i_2=1}^{q^2} \sum_{u_1,\ldots,u_d \ge 0 : u_1+\cdots+u_d \ge 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}), \pi_{j;a_{i_1,j}}(b_{i_1,j,2}), \\
&\qquad \pi_{i_1,j,3}(0), \ldots, \pi_{i_1,j,u_j}(0) : 1 \le j \le d] \\
&\qquad \times \nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}), \pi_{j;a_{i_2,j}}(b_{i_2,j,2}), \pi_{i_2,j,3}(0), \ldots, \pi_{i_2,j,u_j}(0) : 1 \le j \le d]\Big\} \\
&= \frac{1}{q^4} \sum_{i=1}^{q^2} \sum_{u_1,\ldots,u_d \ge 0 : u_1+\cdots+u_d \ge 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \pi_{j;a_{i,j}}(b_{i,j,2}), \\
&\qquad \pi_{i,j,3}(0), \ldots, \pi_{i,j,u_j}(0) : 1 \le j \le d]^2\Big\} \\
&\qquad + \frac{1}{q^4} \sum_{i_1=1}^{q^2} \sum_{i_2 \ne i_1} \sum_{u_1,\ldots,u_d \ge 0 : u_1+\cdots+u_d \ge 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}), \pi_{j;a_{i_1,j}}(b_{i_1,j,2}), \\
&\qquad \pi_{i_1,j,3}(0), \ldots, \pi_{i_1,j,u_j}(0) : 1 \le j \le d] \\
&\qquad \times \nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}), \pi_{j;a_{i_2,j}}(b_{i_2,j,2}), \pi_{i_2,j,3}(0), \ldots, \pi_{i_2,j,u_j}(0) : 1 \le j \le d]\Big\}. \qquad (19)
\end{aligned}$$

Since $A \in \mathrm{OA}(q^2, d, q, 2)$, we have

$$\begin{aligned}
&\frac{1}{q^4} \sum_{i_1=1}^{q^2} \sum_{i_2 \ne i_1} \sum_{u_1,\ldots,u_d \ge 0 : u_1+\cdots+u_d \ge 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}), \pi_{j;a_{i_1,j}}(b_{i_1,j,2}), \\
&\qquad \pi_{i_1,j,3}(0), \ldots, \pi_{i_1,j,u_j}(0) : 1 \le j \le d] \\
&\qquad \times \nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}), \pi_{j;a_{i_2,j}}(b_{i_2,j,2}), \pi_{i_2,j,3}(0), \ldots, \pi_{i_2,j,u_j}(0) : 1 \le j \le d]\Big\} \\
&= \frac{1}{q^4} \sum_{i_1=1}^{q^2} \sum_{0 \le u_1,\ldots,u_d \le 2 : u_1+\cdots+u_d \ge 3} \sum_{i_2 \ne i_1} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}), \pi_{j;a_{i_1,j}}(b_{i_1,j,2}), \\
&\qquad \pi_{i_1,j,3}(0), \ldots, \pi_{i_1,j,u_j}(0) : 1 \le j \le d] \\
&\qquad \times \nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}), \pi_{j;a_{i_2,j}}(b_{i_2,j,2}), \pi_{i_2,j,3}(0), \ldots, \pi_{i_2,j,u_j}(0) : 1 \le j \le d]\Big\} \\
&= \frac{1}{q^4} \sum_{i_1=1}^{q^2} \sum_{0 \le u_1,\cdots,u_d \le 2 : u_1+\cdots+u_d \ge 3} \sum_{i_2 \ne i_1} \sum_{l=1}^{|u|} \mathcal{I}\{a_{i_1,j_l} = a_{i_2,j_l}, u_{j_l} = 1\} \\
&\qquad \times E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}), \pi_{j;a_{i_1,j}}(b_{i_1,j,2}), \pi_{i_1,j,3}(0), \ldots, \pi_{i_1,j,u_j}(0) : 1 \le j \le d]
\end{aligned}$$

$$\begin{aligned}
&\times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\
&+\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l},u_{j_l}=2\}\\
&\times E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\le j\le d]\\
&\quad\times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\
&+\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,j_l}\ne a_{i_2,j_l},\forall l=1,\ldots,|u|\}\\
&\times E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\le j\le d]\\
&\quad\times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}.
\end{aligned}\tag{20}$$

We further note that

$$\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}=\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\text{ and }a_{i_1,j_m}\ne a_{i_2,j_m}\ \forall m\ne l\}.$$

Consequently,

$$\begin{aligned}
&\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l},u_{j_l}=1\}\\
&\times E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\le j\le d]\\
&\quad\times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\
=\;&\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\\
&\quad\times\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\text{ and }u_{j_1}=\cdots=u_{j_{|u|}}=1\}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\}\\
=\;&\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}\\
&\quad\times E\Big\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad\times\Big(\frac{1}{q-1}\Big)^{|u|-1}\Big[\prod_{1\le m\le|u|:m\ne l}\sum_{0\le\tilde c_{j_m}\le q-1:\tilde c_{j_m}\ne\pi_{j_m}(a_{i_1,j_m})}\Big]\\
&\qquad\times\nu^*[\tilde c_{j_1};\ldots;\tilde c_{j_{l-1}};\pi_{j_l}(a_{i_1,j_l});\tilde c_{j_{l+1}};\ldots;\tilde c_{j_{|u|}}]\Big\}
\end{aligned}$$

$$
\begin{aligned}
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\frac{(-1)^{|u|-1}}{(q-1)^{|u|-1}} \\
&\qquad \times\sum_{i_2\neq i_1}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
&= \sum_{0\le u_1,\ldots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|-1}|u|}{q^2(q-1)^{|u|-2}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},
\end{aligned}
$$

$$
\begin{aligned}
&\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\neq i_1}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l},u_{j_l}=2\} \\
&\times E\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\le j\le d] \\
&\qquad \times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\} \\
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\neq i_1} \\
&\qquad \times\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l},u_{j_l}=2 \text{ and } u_{j_k}=1,\forall k\neq l\} \\
&\qquad \times E\Big\{\nu^*_{u_{j_1},\ldots,u_{j_{|u|}}}[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{l-1}}(a_{i_1,j_{l-1}}); \\
&\qquad\quad \pi_{j_l}(a_{i_1,j_l}),\pi_{j_l;a_{i_1,j_l}}(b_{i_1,j_l,2});\pi_{j_{l+1}}(a_{i_1,j_{l+1}});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
&\qquad \times\nu^*_{u_{j_1},\cdots,u_{j_{|u|}}}[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{l-1}}(a_{i_2,j_{l-1}}); \\
&\qquad\quad \pi_{j_l}(a_{i_2,j_l}),\pi_{j_l;a_{i_2,j_l}}(b_{i_2,j_l,2});\pi_{j_{l+1}}(a_{i_2,j_{l+1}});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\neq i_1} \\
&\qquad \times\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l},u_{j_l}=2 \text{ and } u_{j_k}=1,\forall k\neq l\} \\
&\qquad \times E\Big\{\nu^*_{u_{j_1},\ldots,u_{j_{|u|}}}[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{l-1}}(a_{i_1,j_{l-1}}); \\
&\qquad\quad \pi_{j_l}(a_{i_1,j_l}),\pi_{j_l;a_{i_1,j_l}}(b_{i_1,j_l,2});\pi_{j_{l+1}}(a_{i_1,j_{l+1}});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
&\qquad \times(\frac{1}{q-1})^{|u|}\sum_{0\le\tilde{c}_{j_l}\le q-1:\tilde{c}_{j_l}\neq\pi_{j_l;a_{i_1,j_l}}(b_{i_1,j_l,2})} \\
&\qquad \times\Big[\prod_{1\le m\le|u|:m\neq l}\sum_{0\le\tilde{c}_{j_m}\le q-1:\tilde{c}_{j_m}\neq\pi_{j_m}(a_{i_1,j_m})}\Big] \\
&\qquad \times\nu^*_{u_{j_1},\ldots,u_{j_{|u|}}}[\tilde{c}_{j_1};\ldots;\tilde{c}_{j_{l-1}};\pi_{j_l}(a_{i_1,j_l}),\tilde{c}_{j_l};\tilde{c}_{j_{l+1}};\ldots;\tilde{c}_{j_{|u|}}]\Big\}
\end{aligned}
$$

$$
\begin{aligned}
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 2:u_1+\cdots+u_d\ge 3}\frac{(-1)^{|u|}}{(q-1)^{|u|}}\sum_{i_2\ne i_1}\\
&\qquad\times\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}, u_{j_l}=2 \text{ and } u_{j_k}=1, \forall k\ne l\}\\
&\qquad\times E\Big\{\nu^*_{u_{j_1},\dots,u_{j_{|u|}}}[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{l-1}}(a_{i_1,j_{l-1}});\\
&\qquad\quad\pi_{j_l}(a_{i_1,j_l}),\pi_{j_l;a_{i_1,j_l}}(b_{i_1,j_l,2});\pi_{j_{l+1}}(a_{i_1,j_{l+1}});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\Big\}\\
&= \frac{1}{q^2}\sum_{l=1}^{d}\sum_{u_1,\dots,u_d:u_l=2,0\le u_k\le 1\forall k\ne l,|u|+1\ge 3}\frac{(-1)^{|u|}}{(q-1)^{|u|-1}}\\
&\qquad\times E\Big\{\nu^*_{u_{j_1},\dots,u_{j_{|u|}}}[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{l-1}}(a_{1,j_{l-1}});\\
&\qquad\quad\pi_{j_l}(a_{1,j_l}),\pi_{j_l;a_{1,j_l}}(b_{1,j_l,2});\pi_{j_{l+1}}(a_{1,j_{l+1}});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\Big\},
\end{aligned}
$$

and

$$
\begin{aligned}
&\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,j_l}\ne a_{i_2,j_l},\forall l=1,\dots,|u|\}\\
&\times E\Big\{\nu_{u_1,\dots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\dots,\pi_{i_1,j,u_j}(0):1\le j\le d]\\
&\quad\times\nu_{u_1,\dots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\dots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\
=\ &\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 2:u_1+\cdots+u_d\ge 3}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,j_l}\ne a_{i_2,j_l},u_{j_l}=1,\forall l=1,\dots,|u|\}\\
&\times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\dots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\}\\
=\ &\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,j_l}\ne a_{i_2,j_l},\forall l=1,\dots,|u|\}\\
&\times E\Big\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\quad\times(\frac{1}{q-1})^{|u|}\Big[\prod_{1\le m\le|u|}\sum_{0\le\tilde c_{j_m}\le q-1:\tilde c_{j_m}\ne\pi_{j_m}(a_{i_1,j_m})}\Big]\nu^*[\tilde c_{j_1};\dots;\tilde c_{j_{|u|}}]\Big\}\\
=\ &\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|}}{(q-1)^{|u|}}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,j_l}\ne a_{i_2,j_l},\forall l=1,\dots,|u|\}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}\\
=\ &\frac{1}{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|}[q^2-1-|u|(q-1)]}{(q-1)^{|u|}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}
\end{aligned}
$$

$$(21) = \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3} \frac{(-1)^{|u|}(q+1-|u|)}{q^2(q-1)^{|u|-1}} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}.$$

We conclude from (19), (20) and (21) that

$$\begin{aligned}
\sigma^2_{oal} &= \frac{1}{q^2}\sum_{u_1,\dots,u_d\ge 0:u_1+\cdots+u_d\ge 3} E\Big\{\nu_{u_1,\dots,u_d}[\pi_j(a_{1,j}),\pi_{j;a_{1,j}}(b_{1,j,2}),\\
&\qquad \pi_{1,j,3}(0),\dots,\pi_{1,j,u_j}(0):1\le j\le d]^2\Big\}\\
&\quad + \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|-1}|u|}{q^2(q-1)^{|u|-2}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots,\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&\quad + \sum_{l=1}^{d}\sum_{u_1,\dots,u_d:u_l=2,0\le u_k\le 1\forall k\ne l,|u|\ge 2}\frac{(-1)^{|u|}}{q^2(q-1)^{|u|-1}}\\
&\qquad \times E\Big\{\nu^*_{u_{j_1},\dots,u_{j_{|u|}}}[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{l-1}}(a_{1,j_{l-1}});\\
&\qquad\quad \pi_{j_l}(a_{1,j_l}),\pi_{j_l;a_{1,j_l}}(b_{1,j_l,2});\pi_{j_{l+1}}(a_{1,j_{l+1}});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\Big\}\\
&\quad + \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|}(q+1-|u|)}{q^2(q-1)^{|u|-1}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&= \frac{1}{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&\quad + \frac{1}{q^2}\sum_{u_1,\dots,u_d\ge 0:u_1+\cdots+u_d\ge 3\vee(|u|+1)} E\Big\{\nu_{u_1,\dots,u_d}[\pi_j(a_{1,j}),\pi_{j;a_{1,j}}(b_{1,j,2}),\\
&\qquad \pi_{1,j,3}(0),\dots,\pi_{1,j,u_j}(0):1\le j\le d]^2\Big\}\\
&\quad + \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|-1}|u|}{q^2(q-1)^{|u|-2}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots,\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&\quad + \sum_{l=1}^{d}\sum_{u_1,\dots,u_d:u_l=2,0\le u_k\le 1\forall k\ne l,|u|\ge 2}\frac{(-1)^{|u|}}{q^2(q-1)^{|u|-1}}\\
&\qquad \times E\Big\{\nu^*_{u_{j_1},\dots,u_{j_{|u|}}}[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{l-1}}(a_{1,j_{l-1}});\\
&\qquad\quad \pi_{j_l}(a_{1,j_l}),\pi_{j_l;a_{1,j_l}}(b_{1,j_l,2});\pi_{j_{l+1}}(a_{1,j_{l+1}});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\Big\}\\
&\quad + \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|}(q+1-|u|)}{q^2(q-1)^{|u|-1}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}.
\end{aligned}\tag{22}$$

Thus it follows from (22) and Lemma 4 (see Appendix) that

$$\sigma^2_{oal} = \frac{1}{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} + O(\frac{1}{q^3})$$

$$\begin{aligned}
&= \frac{1}{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|\ge 3}\sum_{c_{j_1}=0}^{q-1}\cdots\sum_{c_{j_{|u|}}=0}^{q-1}\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}}\rangle^2 + O(\frac{1}{q^3})\\
&= O(\frac{1}{q^2})\sum_{0\le u_1,\ldots,u_d\le 1:|u|\ge 3} q^{3|u|-3\sum_{l=1}^{|u|}u_{j_l}}\\
&= O(\frac{1}{q^2}),
\end{aligned}$$

as $q\to\infty$. Next we observe from (16) that

$$\begin{aligned}
\sigma_{oas}^2 &= E\Big\{\{\frac{1}{q^2}\sum_{i=1}^{q^2}\sum_{u_1,\ldots,u_d\ge 0:u_1+\cdots+u_d\ge 3}\nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}),\pi_{i,j,2}(0),\ldots,\pi_{i,j,u_j}(0):\\
&\qquad 1\le j\le d]+\frac{1}{q^2}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\nu_2^*[\pi_k(a_{i,k}),\pi_{i,k,2}(0)]\}^2\Big\}\\
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{u_1,\ldots,u_d\ge 0:u_1+\cdots+u_d\ge 3}E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{i_1,j,2}(0),\ldots,\pi_{i_1,j,u_j}(0):\\
&\qquad 1\le j\le d]\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{i_2,j,2}(0),\ldots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\
&\quad +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{k=1}^{d}E\Big\{\nu_2^*[\pi_k(a_{i_1,k}),\pi_{i_1,k,2}(0)]\nu_2^*[\pi_k(a_{i_2,k}),\pi_{i_2,k,2}(0)]\Big\}\\
&= \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{u_1,\ldots,u_d\ge 0:u_1+\cdots+u_d\ge 3}E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{i_1,j,2}(0),\ldots,\pi_{i_1,j,u_j}(0):\\
&\qquad 1\le j\le d]^2\Big\}\\
&\quad +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{k=1}^{d}E\{\nu_2^*[\pi_k(a_{i_1,k}),\pi_{i_1,k,2}(0)]^2\}\\
&\quad +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\ne i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}E\Big\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad \times\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\}\\
&\quad +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\ne i_1}\sum_{k=1}^{d}E\{\nu_2^*[\pi_k(a_{i_1,k}),\pi_{i_1,k,2}(0)]\nu_2^*[\pi_k(a_{i_2,k}),\pi_{i_2,k,2}(0)]\}. \qquad (23)
\end{aligned}$$

From Lemma 4, we obtain

$$\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{k=1}^{d}E\{\nu_2^*[\pi_k(a_{i_1,k}),\pi_{i_1,k,2}(0)]^2\} = O(\frac{1}{q^4}), \qquad (24)$$

$$\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{k=1}^{d} E\{\nu_2^*[\pi_k(a_{i_1,k}),\pi_{i_1,k,2}(0)]\nu_2^*[\pi_k(a_{i_2,k}),\pi_{i_2,k,2}(0)]\} \;\; = \;\; 0,$$

and

$$\begin{aligned}
& \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3} E\Big\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \quad \times\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
= \; & \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\} \\
& \quad \times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\} \\
& +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\mathcal{I}\{a_{i_1,j_l}\neq a_{i_2,j_l}, l=1,\ldots,|u|\} \\
& \quad \times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\} \\
= \; & \frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\sum_{l=1}^{|u|}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\} \\
& \quad \times(-\frac{1}{q-1})^{|u|-1}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
& +\frac{1}{q^4}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:u_1+\cdots+u_d\ge 3}\mathcal{I}\{a_{i_1,j_l}\neq a_{i_2,j_l}, l=1,\ldots,|u|\} \\
& \quad \times(-\frac{1}{q-1})^{|u|}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
= \; & \frac{1}{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|\ge 3}|u|(q-1)(-\frac{1}{q-1})^{|u|-1}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} \\
& +\sum_{0\le u_1,\ldots,u_d\le 1:|u|\ge 3}\frac{(-1)^{|u|}[q^2-1-|u|(q-1)]}{q^2(q-1)^{|u|}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} \\
= \; & O(\frac{1}{q^3}),
\end{aligned}$$

as $q\to\infty$. Hence we conclude from (22) and (23) that

$$\begin{aligned}
\sigma_{oas}^2 \;\; = \;\; & \frac{1}{q^2}\sum_{u_1,\ldots,u_d\ge 0:u_1+\cdots+u_d\ge 3}E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{1,j}),\pi_{1,j,2}(0),\ldots,\pi_{1,j,u_j}(0): \\
& \qquad 1\le j\le d]^2\Big\}+O(\frac{1}{q^3}) \\
= \;\; & \sigma_{oal}^2+O(\frac{1}{q^3}),
\end{aligned}$$

as $q \to \infty$. The remaining case, namely σ^2, can be shown in a similar (though simpler) manner. This proves Proposition 1. $\square$

3 A multivariate central limit theorem

First we define the proxy statistic

$$W = \frac{1}{q^2\sigma}\sum_{i=1}^{q^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|\ge 3} \nu^*[\pi_{j_1}(a_{i,j_1});\dots;\pi_{j_{|u|}}(a_{i,j_{|u|}})], \tag{25}$$

where σ^2 is as in (18). Clearly we have $E(W^2)=1$.

Proposition 2. *Let $f:[0,1)^d\to\mathbb{R}$ be smooth with a Lipschitz continuous mixed partial of order d. Suppose further that W_{oal}, W_{oas} and W are as defined by (15), (16) and (25) respectively with $A\in OA(q^2,d,q,2)$. Then $q(\sigma_{oal}W_{oal}-\sigma W)\to 0$ and $q(\sigma_{oas}W_{oas}-\sigma W)\to 0$ in probability as $q\to\infty$.*

Proof. We observe that

$$q(\sigma_{oal}W_{oal}-\sigma W)=\sum_{r=1}^{d}\sum_{1\le k_1<\cdots<k_r\le d}\Delta_{k_1,\dots,k_r},$$

where

$$\begin{aligned}\Delta_{k_1,\dots,k_r} &= \frac{1}{q}\sum_{i=1}^{q^2}\sum_{u_1,\dots,u_d\ge 0:u_k\ge 2\Leftrightarrow k\in\{k_1,\dots,k_r\},u_1+\cdots+u_d\ge 3} \nu_{u_1,\dots,u_d}[\pi_j(a_{i,j}),\\ &\qquad \pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\dots,\pi_{i,j,u_j}(0):1\le j\le d].\end{aligned}$$

Here $\Leftrightarrow$ denotes if and only if and that given $u_1,\dots,u_d\ge 0$, we write $u_k\ge 1\Leftrightarrow k\in\{j_1,\dots,j_{|u|}\}$ where $|u|=\sum_{i=1}^d \mathcal{I}\{u_i\ge 1\}$. Now for $r=1,\dots,d$, we have

$$\begin{aligned}E(\Delta^2_{1,\dots,r}) &= \frac{1}{q^2}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{u_1,\dots,u_d\ge 0:u_k\ge 2\Leftrightarrow k\in\{1,\dots,r\},u_1+\cdots+u_d\ge 3} E\Big\{\nu_{u_1,\dots,u_d}[\pi_j(a_{i_1,j}),\\ &\qquad \pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\dots,\pi_{i_1,j,u_j}(0):1\le j\le d]\\ &\qquad \times\nu_{u_1,\dots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\dots,\pi_{i_2,j,u_j}(0):1\le j\le d]\Big\}\\ &= \frac{1}{q^2}\sum_{i=1}^{q^2}\sum_{u_1,\dots,u_d\ge 0:u_k\ge 2\Leftrightarrow k\in\{1,\dots,r\},u_1+\cdots+u_d\ge 3} E\Big\{\nu_{u_1,\dots,u_d}[\pi_j(a_{i,j}),\\ &\qquad \pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\dots,\pi_{i,j,u_j}(0):1\le j\le d]^2\Big\}\end{aligned}$$

$$
\begin{aligned}
&+\frac{1}{q^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{u_1,\ldots,u_d\geq 0:u_k\geq 2\Leftrightarrow k\in\{1,\ldots,r\},u_1+\cdots+u_d\geq 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\\
&\quad \pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\leq j\leq d]\\
&\quad \times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\leq j\leq d]\Big\}. \qquad (26)
\end{aligned}
$$

Using the fact that $A\in\mathrm{OA}(q^2,d,q,2)$, we further observe that

$$
\begin{aligned}
&\frac{1}{q^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{u_1,\ldots,u_d\geq 0:u_k\geq 2\Leftrightarrow k\in\{1,\ldots,r\},u_1+\cdots+u_d\geq 3} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\\
&\quad \pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\leq j\leq d]\\
&\quad \times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\leq j\leq d]\Big\}\\
={}&\frac{1}{q^2}\sum_{i_1=1}^{q^2}\sum_{0\leq u_1,\ldots,u_d\leq 2:u_k=2\Leftrightarrow k\in\{1,\ldots,r\},u_1+\cdots+u_d\geq 3}\sum_{i_2\neq i_1} E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\\
&\quad \pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\leq j\leq d]\\
&\quad \times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\leq j\leq d]\Big\}\\
={}&\frac{\mathcal{I}\{r=1\}}{q^2}\sum_{i_1=1}^{q^2}\sum_{0\leq u_1,\ldots,u_d\leq 2:u_k=2\Leftrightarrow k=1,|u|+1\geq 3}\sum_{i_2\neq i_1}\mathcal{I}\{a_{i_1,1}=a_{i_2,1}\}\\
&\quad \times E\Big\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_1,j}),\pi_{j;a_{i_1,j}}(b_{i_1,j,2}),\pi_{i_1,j,3}(0),\ldots,\pi_{i_1,j,u_j}(0):1\leq j\leq d]\\
&\quad \times\nu_{u_1,\ldots,u_d}[\pi_j(a_{i_2,j}),\pi_{j;a_{i_2,j}}(b_{i_2,j,2}),\pi_{i_2,j,3}(0),\ldots,\pi_{i_2,j,u_j}(0):1\leq j\leq d]\Big\}\\
={}&\frac{\mathcal{I}\{r=1\}}{q^2}\sum_{i_1=1}^{q^2}\sum_{0\leq u_1,\ldots,u_d\leq 2:u_k=2\Leftrightarrow k=1,|u|\geq 2}\sum_{i_2\neq i_1}\mathcal{I}\{a_{i_1,1}=a_{i_2,1}\}\\
&\quad \times E\Big\{\nu^*_{u_1,u_{j_2},\ldots,u_{j_{|u|}}}[\pi_1(a_{i_1,1}),\pi_{1;a_{i_1,1}}(b_{i_1,1,2});\pi_{j_2}(a_{i_1,j_2});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\quad \times\nu^*_{u_1,u_{j_2},\ldots,u_{j_{|u|}}}[\pi_1(a_{i_2,1}),\pi_{1;a_{i_2,1}}(b_{i_2,1,2});\pi_{j_2}(a_{i_2,j_2});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\}\\
={}&\frac{\mathcal{I}\{r=1\}}{q^2}\sum_{i_1=1}^{q^2}\sum_{0\leq u_1,\ldots,u_d\leq 2:u_k=2\Leftrightarrow k=1,|u|\geq 2}\sum_{i_2\neq i_1}\mathcal{I}\{a_{i_1,1}=a_{i_2,1}\}\\
&\quad \times E\Big\{\nu^*_{u_1,u_{j_2},\ldots,u_{j_{|u|}}}[\pi_1(a_{i_1,1}),\pi_{1;a_{i_1,1}}(b_{i_1,1,2});\pi_{j_2}(a_{i_1,j_2});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\quad \times(\frac{1}{q-1})^{|u|}\sum_{0\leq\tilde{c}_1\leq q-1:\tilde{c}_1\neq\pi_{1;a_{i_1,1}}(b_{i_1,1,2})}\Big[\prod_{2\leq m\leq|u|}\sum_{0\leq\tilde{c}_{j_m}\leq q-1:\tilde{c}_{j_m}\neq\pi_{j_m}(a_{i_1,j_m})}\Big]\\
&\quad \times\nu^*_{u_1,u_{j_2},\ldots,u_{j_{|u|}}}[\pi_1(a_{i_1,1}),\tilde{c}_1;\tilde{c}_{j_2};\ldots;\tilde{c}_{j_{|u|}}]\Big\}
\end{aligned}
$$

$$\begin{aligned}
&= \frac{\mathcal{I}\{r=1\}}{q^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 2: u_k=2\Leftrightarrow k=1,|u|\ge 2}\frac{(-1)^{|u|}}{(q-1)^{|u|}}\sum_{i_2\ne i_1}\mathcal{I}\{a_{i_1,1}=a_{i_2,1}\}\\
&\quad\times E\{\nu^*_{u_1,u_{j_2},\dots,u_{j_{|u|}}}[\pi_1(a_{i_1,1}),\pi_{1;a_{i_1,1}}(b_{i_1,1,2});\pi_{j_2}(a_{i_1,j_2});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}\\
&= \mathcal{I}\{r=1\}\sum_{0\le u_1,\dots,u_d\le 2: u_k=2\Leftrightarrow k=1,|u|\ge 2}\frac{(-1)^{|u|}}{(q-1)^{|u|-1}}\\
&\quad\times E\{\nu^*_{u_1,u_{j_2},\dots,u_{j_{|u|}}}[\pi_1(a_{1,1}),\pi_{1;a_{1,1}}(b_{1,1,2});\pi_{j_2}(a_{1,j_2});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}. \qquad (27)
\end{aligned}$$

It follows from (26) and (27) that

$$\begin{aligned}
&q^2E[(\sigma_{oal}W_{oal}-\sigma W)^2]\\
&= E[(\sum_{r=1}^{d}\sum_{1\le j_{k_1}<\cdots<j_{k_r}\le d}\Delta_{j_{k_1},\dots,j_{k_r}})^2]\\
&= \sum_{r=1}^{d}\sum_{1\le j_{k_1}<\cdots<j_{k_r}\le d}E(\Delta^2_{j_{k_1},\dots,j_{k_r}})\\
&= \sum_{r=1}^{d}\sum_{1\le j_{k_1}<\cdots<j_{k_r}\le d}\sum_{u_1,\dots,u_d\ge 0: u_k\ge 2\Leftrightarrow k\in\{j_{k_1},\dots,j_{k_r}\},u_1+\cdots+u_d\ge 3}\\
&\qquad\times E\{\nu_{u_1,\dots,u_d}[\pi_j(a_{1,j}),\pi_{j;a_{1,j}}(b_{1,j,2}),\pi_{1,j,3}(0),\dots,\pi_{1,j,u_j}(0):1\le j\le d]^2\}\\
&\quad+\sum_{r=1}^{d}\sum_{0\le u_1,\dots,u_d\le 2: u_k=2\Leftrightarrow k=r,|u|\ge 2}\frac{(-1)^{|u|}}{(q-1)^{|u|-1}}\\
&\qquad\times E\{\nu_{u_1,\dots,u_d}[\pi_j(a_{1,j}),\pi_{j;a_{1,j}}(b_{1,j,2}),\pi_{1,j,3}(0),\dots,\pi_{1,j,u_j}(0):1\le j\le d]^2\}. \qquad (28)
\end{aligned}$$

Finally from (28) and Lemma 4 (see Appendix), we have

$$\begin{aligned}
&q^2E[(\sigma_{oal}W_{oal}-\sigma W)^2]\\
&= \sum_{r=1}^{d}\sum_{1\le j_{k_1}<\cdots<j_{k_r}\le d}\sum_{u_1,\dots,u_d\ge 0: u_k\ge 2\Leftrightarrow k\in\{j_{k_1},\dots,j_{k_r}\},u_1+\cdots+u_d\ge 3}\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\cdots\\
&\qquad\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2\sum_{\alpha=0}^{|u|}\binom{|u|}{\alpha}\frac{1}{q^\alpha}(1-\frac{1}{q})^{|u|-\alpha}\\
&\quad+\sum_{r=1}^{d}\sum_{0\le u_1,\dots,u_d\le 2: u_k=2\Leftrightarrow k=r,|u|\ge 2}\frac{(-1)^{|u|}}{(q-1)^{|u|-1}}\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\cdots\\
&\qquad\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2\sum_{\alpha=0}^{|u|}\binom{|u|}{\alpha}\frac{1}{q^\alpha}(1-\frac{1}{q})^{|u|-\alpha}
\end{aligned}$$

$$\begin{aligned} &= \sum_{r=1}^{d} \binom{d}{r} \sum_{u_1,\ldots,u_d \geq 0: u_k \geq 2 \Leftrightarrow k \in \{1,\ldots,r\}, u_1+\cdots+u_d \geq 3} O(q^{2|u|-2(u_1+\cdots+u_d)}) \\ &\quad + \sum_{0 \leq u_1,\ldots,u_d \leq 2: u_k = 2 \Leftrightarrow k=1, |u| \geq 2} \frac{(-1)^{|u|} d}{(q-1)^{|u|-1}} O(q^{2|u|-2(u_1+\cdots+u_d)}) \\ &= O(q^{-2}), \end{aligned}$$

as $q \to \infty$. Using Chebyshev's inequality, we conclude that $q(\sigma_{oal} W_{oal} - \sigma W) \to 0$ in probability as $q \to \infty$. Next we observe from (24) that

$$E\Big\{\{\frac{1}{q}\sum_{i=1}^{q^2}\sum_{k=1}^{d} \nu_2^*[\pi_k(a_{i,k}), \pi_{i,k,2}(0)]\}^2\Big\} = O(\frac{1}{q^2}),$$

as $q \to \infty$. Again by Chebyshev's inequality, we have $q^{-1}\sum_{i=1}^{q^2}\sum_{k=1}^{d} \nu_2^*[\pi_k(a_{i,k}), \pi_{i,k,2}(0)] \to 0$ in probability as $q \to \infty$. Thus we conclude from (16) that to prove that $q(\sigma_{oas} W_{oas} - \sigma W) \to 0$ in probability as $q \to \infty$, it suffices to show that

$$\begin{aligned} &\frac{1}{q}\sum_{i=1}^{q^2} \sum_{u_1,\ldots,u_d \geq 0: u_1+\cdots+u_d \geq 3} \nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}), \pi_{i,j,2}(0), \ldots, \pi_{i,j,u_j}(0) : 1 \leq j \leq d] - q\sigma W \\ \to\ & 0, \end{aligned}$$

in probability as $q \to \infty$. The proof of the latter statement is similar to the proof that $q(\sigma_{oal} W_{oal} - \sigma W) \to 0$ and hence will be omitted. This proves Proposition 2. □

REMARK. We observe from Theorem 1 and Proposition 1 that for $\int_{[0,1)^d} f_{rem}^2(x)dx > 0$, we have

$$\frac{\sigma_{oal}^2}{\sigma^2} = 1 + O(\frac{1}{q}), \quad \frac{\sigma_{oas}^2}{\sigma^2} = 1 + O(\frac{1}{q}), \quad \text{as } q \to \infty.$$

Thus we conclude from Proposition 2 that to show W_{oal} and W_{oas} both tend in law to the standard (univariate) normal distribution as $q \to \infty$, it suffices to show that the proxy statistic W tends in law to that distribution.

Suppose $d \geq 3$. For $\ell = 1, \ldots, d-2$, we define

$$\begin{aligned} \sigma_\ell^2 &= E\Big\{\{\frac{1}{q^2}\sum_{i=1}^{q^2} \sum_{0 \leq u_1,\ldots,u_d \leq 1: |u| = \ell+2} \nu^*[\pi_{j_1}(a_{i,j_1}); \ldots; \pi_{j_{|u|}}(a_{i,j_{|u|}})]\}^2\Big\}, \\ V_\ell &= \frac{1}{q^2 \sigma_\ell}\sum_{i=1}^{q^2} \sum_{0 \leq u_1,\ldots,u_d \leq 1: |u| = \ell+2} \nu^*[\pi_{j_1}(a_{i,j_1}); \ldots; \pi_{j_{|u|}}(a_{i,j_{|u|}})], \end{aligned}$$

$$\text{and } V = (V_1, \ldots, V_{d-2})' \text{ where } |u| = \textstyle\sum_{i=1}^{d} \mathcal{I}\{u_i \geq 1\}. \tag{29}$$

We shall prove that the random vector V converges weakly to the standard $(d-2)$-variate normal distribution Φ_{d-2} as q tends to infinity. To do so, we shall use the multivariate normal version of Stein's method [see Stein (1972), (1986)] as given in Götze (1991) and Bolthausen and Götze (1993).

Let $\mathcal{A}$ be a class of measurable functions from $\mathbb{R}^{d-2} \to \mathbb{R}$ such that $\sup_{v\in\mathbb{R}^{d-2}} |g(v)| \leq 1$ for all $g \in \mathcal{A}$. For $g \in \mathcal{A}$ and $\delta > 0$, define

$$\begin{aligned} g_\delta^+(v) &= \sup\{g(v+y) : \|y\| \leq \delta\}, \quad \forall v \in \mathbb{R}^{d-2}, \\ g_\delta^-(v) &= \inf\{g(v+y) : \|y\| \leq \delta\}, \quad \forall v \in \mathbb{R}^{d-2}, \\ \omega(g,\delta) &= \int_{\mathbb{R}^{d-2}} [g_\delta^+(y) - g_\delta^-(y)]\Phi_{d-2}(dy). \end{aligned}$$

We further assume that $\mathcal{A}$ is closed under supremum and affine transformations, that is, $g \in \mathcal{A}$ implies that $g_\delta^+ \in \mathcal{A}$, $g_\delta^- \in \mathcal{A}$ and $g \circ T \in \mathcal{A}$ whenever $T : \mathbb{R}^{d-2} \to \mathbb{R}^{d-2}$ is affine. Finally we assume that there exists a constant $\Delta \geq 2\sqrt{d-2}$ such that

$$\sup\{\omega(g,\delta) : g \in \mathcal{A}\} \leq \Delta\delta, \quad \forall \delta > 0. \tag{30}$$

We observe from Bolthausen and Götze (1993) that $\mathcal{A}$ can be taken to be the class of all indicator functions of measurable convex sets in $\mathbb{R}^{d-2}$. For $h \in \mathcal{A}$ and $0 \leq t \leq 1$, define

$$\begin{aligned} \chi_t(v|h) &= \int_{\mathbb{R}^{d-2}} [h(y) - h(t^{1/2}y + (1-t)^{1/2}v)]\Phi_{d-2}(dy), \qquad (31) \\ \psi_t(v) &= \frac{1}{2}\int_t^1 \chi_s(v|h)\frac{ds}{1-s}, \quad \forall v \in \mathbb{R}^{d-2}. \end{aligned}$$

Then $-\chi_0(v|h) = h(v) - \Phi_{d-2}(h)$ where $\Phi_{d-2}(h) = Eh(Z)$ and Z is a random vector having distribution Φ_{d-2}. The following two lemmas are due to Götze (1991). Since the proofs are only briefly sketched in Götze (1991), detailed proofs of Lemmas 1 and 2 are given below.

Lemma 1. *For* $0 < t < 1$ *and* $v = (v_1, \ldots, v_{d-2})' \in \mathbb{R}^{d-2}$, *we have*

$$\sum_{i=1}^{d-2} \frac{\partial^2}{\partial v_i^2}\psi_t(v) - \sum_{i=1}^{d-2} v_i \frac{\partial}{\partial v_i}\psi_t(v) = -\chi_t(v|h). \tag{32}$$

There exists a constant c (depending only on d) such that

$$\sup_{1\leq i,j\leq d-2}\ \sup_{v\in\mathbb{R}^{d-2}} |\frac{\partial^2}{\partial v_i \partial v_j}\psi_t(v)| \leq \|h\|_\infty \log(1/t),$$

where $\|h\|_\infty = \sup_{v\in\mathbb{R}^{d-2}} |h(v)|$ *and*

$$\begin{aligned} &\sup_{1\leq i,j,k\leq d-2} |\int_{\mathbb{R}^{d-2}} [\frac{\partial^3}{\partial v_i \partial v_j \partial v_k}\psi_t(v)]Q(dv)| \\ \leq\ & \frac{c}{t^{1/2}} \sup\{|\int_{\mathbb{R}^{d-2}} h(sv+y)Q(dv)| : 0 \leq s \leq 1, y \in \mathbb{R}^{d-2}\} \end{aligned}$$

for all finite signed measures Q *on* $\mathbb{R}^{d-2}$.

PROOF. We first observe from (31) that

$$\chi_t(v|h) = \int_{\mathbb{R}^{d-2}} h(y)\Phi_{d-2}(dy) - \int_{\mathbb{R}^{d-2}} \frac{h(y)}{(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy.$$

Now we consider the partial differential equation

$$2(1-t)\frac{\partial}{\partial t}u_t(v) = \sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}u_t(v) - \sum_{i=1}^{d-2} v_i \frac{\partial}{\partial v_i}u_t(v), \qquad \forall t > 0. \tag{33}$$

We observe that a trivial solution to (33) is $u_t(v) = \int_{\mathbb{R}^{d-2}} h(y)\Phi_{d-2}(dy)$. Writing

$$\tilde{u}_{y,t}(v) = \frac{1}{(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)},$$

we have

$$\begin{aligned}
\sum_{i=1}^{d-2} v_i \frac{\partial}{\partial v_i}\tilde{u}_{y,t}(v) &= \frac{(1-t)^{1/2}}{t(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)} \sum_{i=1}^{d-2} v_i(y_i-(1-t)^{1/2}v_i),\\
\sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\tilde{u}_{y,t}(v) &= \frac{1-t}{t^2(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)} \sum_{i=1}^{d-2}[(y_i-(1-t)^{1/2}v_i)^2 - t]\\
&= \frac{1-t}{t^2(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)} \sum_{i=1}^{d-2}(y_i-(1-t)^{1/2}v_i)^2\\
&\quad -\frac{(d-2)(1-t)}{t(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)},
\end{aligned}$$

and

$$\begin{aligned}
2(1-t)\frac{\partial}{\partial t}\tilde{u}_{y,t}(v) &= -\frac{(d-2)(1-t)}{t(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}\\
&\quad -\frac{1-t}{(2\pi t)^{(d-2)/2}} e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}\\
&\qquad \times \sum_{i=1}^{d-2}[-\frac{(y_i-(1-t)^{1/2}v_i)^2}{t^2} + \frac{v_i(y_i-(1-t)^{1/2}v_i)}{t(1-t)^{1/2}}].
\end{aligned}$$

Thus $\tilde{u}_{y,t}(v)$ is a solution to (33) too for all $y \in \mathbb{R}^{d-2}$. Using the dominated convergence theorem, we have

$$\begin{aligned}
2(1-t)\frac{\partial}{\partial t}\chi_t(v|h) &= -2(1-t)\frac{\partial}{\partial t}\int_{\mathbb{R}^{d-2}} h(y)\tilde{u}_{y,t}(v)dy\\
&= -2(1-t)\int_{\mathbb{R}^{d-2}} h(y)\frac{\partial}{\partial t}\tilde{u}_{y,t}(v)dy
\end{aligned}$$

$$\begin{aligned} &= \sum_{i=1}^{d-2}\int_{\mathbb{R}^{d-2}} h(y)v_i\frac{\partial}{\partial v_i}\tilde{u}_{y,t}(v)dy - \sum_{i=1}^{d-2}\int_{\mathbb{R}^{d-2}} h(y)\frac{\partial^2}{\partial v_i^2}\tilde{u}_{y,t}(v)dy \\ &= \sum_{i=1}^{d-2} v_i\frac{\partial}{\partial v_i}\int_{\mathbb{R}^{d-2}} h(y)\tilde{u}_{y,t}(v)dy - \sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\int_{\mathbb{R}^{d-2}} h(y)\tilde{u}_{y,t}(v)dy \end{aligned}$$

$$= \sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\chi_t(v|h) - \sum_{i=1}^{d-2} v_i\frac{\partial}{\partial v_i}\chi_t(v|h). \tag{34}$$

We further observe that

$$\begin{aligned} \frac{\partial}{\partial v_i}\chi_t(v|h) &= -\frac{\partial}{\partial v_i}\int_{\mathbb{R}^{d-2}}\frac{h(y)}{(2\pi t)^{(d-2)/2}}e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy \\ &= -\frac{(1-t)^{1/2}}{t(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy, \\ \frac{\partial^2}{\partial v_i^2}\chi_t(v|h) &= -\frac{(1-t)^{1/2}}{t(2\pi t)^{(d-2)/2}}\frac{\partial}{\partial v_i}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy \\ &= \frac{1-t}{t(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy \\ &\quad -\frac{1-t}{t^2(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)^2e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy, \\ \frac{\partial^3}{\partial v_i^3}\chi_t(v|h) &= \frac{3(1-t)^{3/2}}{t^2(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy \\ &\quad -\frac{(1-t)^{3/2}}{t^3(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)^3e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy. \end{aligned}$$

For $1 \le i \ne j \le d-2$, we have

$$\begin{aligned} \frac{\partial^3}{\partial v_i^2\partial v_j}\chi_t(v|h) &= \frac{(1-t)^{3/2}}{t^2(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_j-(1-t)^{1/2}v_j)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy \\ &\quad -\frac{(1-t)^{3/2}}{t^3(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)^2(y_j-(1-t)^{1/2}v_j) \\ &\quad \times e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy, \end{aligned}$$

and for i, j, k all distinct, we have

$$\begin{aligned} \frac{\partial^3}{\partial v_i\partial v_j\partial v_k}\chi_t(v|h) &= -\frac{(1-t)^{3/2}}{t^3(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-t)^{1/2}v_i)(y_j-(1-t)^{1/2}v_j) \\ &\quad \times(y_k-(1-t)^{1/2}v_k)e^{-\sum_{l=1}^{d-2}(y_l-(1-t)^{1/2}v_l)^2/(2t)}dy. \end{aligned}$$

Consequently,

$$|\frac{\partial}{\partial v_i}\chi_t(v|h)| \le \frac{(1-t)^{1/2}\|h\|_\infty}{t(2\pi t)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}}|y_i|e^{-\sum_{l=1}^{d-2}y_l^2/(2t)}dy = \frac{(1-t)^{1/2}\|h\|_\infty}{t^{1/2}},$$

$$|\frac{\partial^2}{\partial v_i^2}\chi_t(v|h)| \le \frac{2(1-t)\|h\|_\infty}{t}. \tag{35}$$

Since $\chi_1(v|h) = 0$, it follows from (34) that

$$\begin{aligned}
-\chi_t(v|h) &= \chi_1(v|h) - \chi_t(v|h) \\
&= \int_t^1 [\frac{\partial}{\partial s}\chi_s(v|h)]ds \\
&= \frac{1}{2}\int_t^1 \sum_{i=1}^{d-2} \frac{\partial^2}{\partial v_i^2}\chi_s(v|h)\frac{ds}{1-s} - \frac{1}{2}\int_t^1 \sum_{i=1}^{d-2} v_i \frac{\partial}{\partial v_i}\chi_s(v|h)\frac{ds}{1-s} \\
&= \sum_{i=1}^{d-2} \frac{\partial^2}{\partial v_i^2}\psi_t(v) - \sum_{i=1}^{d-2} v_i \frac{\partial}{\partial v_i}\psi_t(v), \qquad \forall 0 < t < 1.
\end{aligned}$$

The interchange of integration and partial differentiation is justified via (35) and the dominated convergence theorem. This proves (32). Now we observe that for $v \in \mathbb{R}^{d-2}$,

$$\begin{aligned}
\frac{\partial^2}{\partial v_i^2}\psi_t(v) &= \frac{1}{2}\int_t^1 [\frac{\partial^2}{\partial v_i^2}\chi_s(v|h)]\frac{ds}{1-s} \\
&= \frac{1}{2}\int_t^1 \frac{ds}{s(2\pi s)^{(d-2)/2}} \int_{\mathbb{R}^{d-2}} h(y) e^{-\sum_{l=1}^{d-2}[y_l-(1-s)^{1/2}v_l]^2/(2s)} \\
&\quad \times\{1 - \frac{[y_i-(1-s)^{1/2}v_i]^2}{s}\}dy, \\
\frac{\partial^2}{\partial v_i \partial v_j}\psi_t(v) &= \frac{1}{2}\int_t^1 \frac{ds}{s(2\pi s)^{(d-2)/2}} \int_{\mathbb{R}^{d-2}} h(y) e^{-\sum_{l=1}^{d-2}[y_l-(1-s)^{1/2}v_l]^2/(2s)} \\
&\quad \times \frac{[y_i-(1-s)^{1/2}v_i][y_j-(1-s)^{1/2}v_j]}{s}dy, \qquad \text{if } 1 \le i \ne j \le d-2,
\end{aligned}$$

and hence

$$\sup_{1\le i,j\le d-2}\ \sup_{v\in\mathbb{R}^{d-2}} |\frac{\partial^2}{\partial v_i \partial v_j}\psi_t(v)| \le \|h\|_\infty \log(1/t).$$

Next we observe that

$$\begin{aligned}
\int_{\mathbb{R}^{d-2}} [\frac{\partial^3}{\partial v_i^3}\psi_t(v)]Q(dv) &= \frac{1}{2}\int_{\mathbb{R}^{d-2}} Q(dv) \frac{\partial^3}{\partial v_i^3}\int_t^1 \chi_s(v|h)\frac{ds}{1-s} \\
&= \frac{1}{2}\int_{\mathbb{R}^{d-2}} Q(dv) \int_t^1 \frac{(1-s)^{1/2}ds}{s^2(2\pi s)^{(d-2)/2}} \\
&\quad \times\Big[3\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-s)^{1/2}v_i)e^{-\|y-(1-s)^{1/2}v\|^2/(2s)}dy \\
&\qquad -\frac{1}{s}\int_{\mathbb{R}^{d-2}} h(y)(y_i-(1-s)^{1/2}v_i)^3 e^{-\|y-(1-s)^{1/2}v\|^2/(2s)}dy\Big] \\
&= \frac{1}{2}\int_t^1 \frac{(1-s)^{1/2}ds}{s^2(2\pi s)^{(d-2)/2}} \int_{\mathbb{R}^{d-2}} \Big[\int_{\mathbb{R}^{d-2}} h(y+(1-s)^{1/2}v)Q(dv)\Big] \\
&\quad \times\Big[3y_i e^{-\|y\|^2/(2s)} - \frac{1}{s}y_i^3 e^{-\|y\|^2/(2s)}\Big]dy,
\end{aligned}$$

where $\|.\|$ is the usual Euclidean norm in $\mathbb{R}^{d-2}$ and

$$\begin{aligned}|\int_{\mathbb{R}^{d-2}}[\frac{\partial^3}{\partial v_i^3}\psi_t(v)]Q(dv)| &\leq \frac{1}{2}\sup\{|\int_{\mathbb{R}^{d-2}}h(y+sv)Q(dv)|: 0\leq s\leq 1, y\in\mathbb{R}^{d-2}\}\\ &\quad\times\int_t^1\frac{(1-s)^{1/2}ds}{s^{3/2}(2\pi)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}}(3|y_i|+|y_i^3|)e^{-\|y\|^2/2}dy\\ &\leq \frac{c}{t^{1/2}}\sup\{|\int_{\mathbb{R}^{d-2}}h(y+sv)Q(dv)|: s\in[0.1], y\in\mathbb{R}^{d-2}\}.\end{aligned}$$

In a similar way, we have for i, j, k all distinct,

$$\begin{aligned}\int_{\mathbb{R}^{d-2}}[\frac{\partial^3}{\partial v_i^2\partial v_j}\psi_t(v)]Q(dv) &= \frac{1}{2}\int_{\mathbb{R}^{d-2}}Q(dv)\int_t^1\frac{(1-s)^{1/2}ds}{s^2(2\pi s)^{(d-2)/2}}\\ &\quad\times\int_{\mathbb{R}^{d-2}}h(y+(1-s)^{1/2}v)(y_j-\frac{y_i^2y_j}{s})e^{-\|y\|^2/(2s)}dy,\\ \int_{\mathbb{R}^{d-2}}[\frac{\partial^3}{\partial v_i\partial v_j\partial v_k}\psi_t(v)]Q(dv) &= \frac{1}{2}\int_{\mathbb{R}^{d-2}}Q(dv)\int_t^1\frac{(1-s)^{1/2}ds}{s^3(2\pi s)^{(d-2)/2}}\\ &\quad\times\int_{\mathbb{R}^{d-2}}h(y+(1-s)^{1/2}v)y_iy_jy_ke^{-\|y\|^2/(2s)}dy,\end{aligned}$$

and

$$\begin{aligned}|\int_{\mathbb{R}^{d-2}}[\frac{\partial^3}{\partial v_i^2\partial v_j}\psi_t(v)]Q(dv)| &\leq \frac{1}{2}\sup\{|\int_{\mathbb{R}^{d-2}}h(y+sv)Q(dv)|: s\in[0,1], y\in\mathbb{R}^{d-2}\}\\ &\quad\times\int_t^1\frac{(1-s)^{1/2}ds}{s^{3/2}(2\pi)^{(d-2)/2}}\int_{\mathbb{R}^{d-2}}(|y_j|+|y_i^2y_j|)e^{-\|y\|^2/2}dy\\ &\leq \frac{c}{t^{1/2}}\sup\{|\int_{\mathbb{R}^{d-2}}h(y+sv)Q(dv)|: s\in[0,1], y\in\mathbb{R}^{d-2}\},\\ |\int_{\mathbb{R}^{d-2}}[\frac{\partial^3}{\partial v_i\partial v_j\partial v_k}\psi_t(v)]Q(dv)| &\leq \frac{c}{t^{1/2}}\sup\{|\int_{\mathbb{R}^{d-2}}h(y+sv)Q(dv)|: s\in[0,1], y\in\mathbb{R}^{d-2}\}.\end{aligned}$$

This proves Lemma 1. □

Lemma 2. *Suppose that (30) holds. Let $0<\varepsilon<1/2$ and Q be a probability distribution on $\mathbb{R}^{d-2}$. Then there exists a constant c_d depending only on d such that*

$$\sup_{g\in\mathcal{A}}|\int_{\mathbb{R}^{d-2}}g(v)[Q(dv)-\Phi_{d-2}(dv)]|\leq c_d[\sup_{h\in\mathcal{A}}|\int_{\mathbb{R}^{d-2}}\chi_{\varepsilon^2}(v|h)Q(dv)|+\Delta\varepsilon].$$

PROOF. Let $m_d\in\mathbb{Z}^+$ and $\alpha>1/2$ be constants satisfying

$$\Phi_{d-2}(\{\|x\|\leq\sqrt{3m_d/4}: x\in\mathbb{R}^{d-2}\})=\alpha.$$

We further write $b=(1-\varepsilon^2)^{1/2}$, $\delta=\varepsilon/b$ and $\Phi_{d-2,\varepsilon}(A)=\Phi_{d-2}(A/\varepsilon), Q_\varepsilon(A)=Q(A/\varepsilon)$ for all Borel sets $A\subset\mathbb{R}^{d-2}$. For $g\in\mathcal{A}$, we define

$$\gamma^*(g;\varepsilon) = \sup\Big\{\max\{\int_{\mathbb{R}^{d-2}}g_\varepsilon^+(v+y)(Q-\Phi_{d-2})*\Phi_{d-2,\varepsilon/\sqrt{m_d(1-\varepsilon^2)}}(dv),$$

$$
\begin{aligned}
& -\int_{\mathbb{R}^{d-2}} g_\varepsilon^-(v+y)(Q-\Phi_{d-2}) * \Phi_{d-2,\varepsilon/\sqrt{m_d(1-\varepsilon^2)}}(dv)\} : y \in \mathbb{R}^{d-2}\Big\}, \\
\tau^*(g;\varepsilon) \quad = \quad & \sup\Big\{ \max\{\int_{\mathbb{R}^{d-2}} [g_\varepsilon^+(v+y) - g(v+y)]\Phi_{d-2}(dv), \\
& -\int_{\mathbb{R}^{d-2}} [g_\varepsilon^-(v+y) - g(v+y)]\Phi_{d-2}(dv)\} : y \in \mathbb{R}^{d-2}\Big\}.
\end{aligned}
$$

Since $\mathcal{A}$ is closed under affine transformations, it follows from (30) that $\tau^*(g;\varepsilon) \le \Delta\varepsilon$. We further observe that

$$
\begin{aligned}
& \sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} h(v)(Q-\Phi_{d-2}) * \Phi_{d-2,\varepsilon/\sqrt{1-\varepsilon^2}}(dv)| \\
= \quad & \sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}}\int_{\mathbb{R}^{d-2}} h(v+\frac{\varepsilon y}{\sqrt{1-\varepsilon^2}})Q(dv)\Phi_{d-2}(dy) \\
& -\int_{\mathbb{R}^{d-2}}\int_{\mathbb{R}^{d-2}} h(v+\frac{\varepsilon y}{\sqrt{1-\varepsilon^2}})\Phi_{d-2}(dv)\Phi_{d-2}(dy)| \\
= \quad & \sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}}\int_{\mathbb{R}^{d-2}} h((1-\varepsilon^2)^{1/2}v+\varepsilon y)Q(dv)\Phi_{d-2}(dy) \\
& -\int_{\mathbb{R}^{d-2}}\int_{\mathbb{R}^{d-2}} h((1-\varepsilon^2)^{1/2}v+\varepsilon y)\Phi_{d-2}(dv)\Phi_{d-2}(dy)| \\
= \quad & \sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}}\int_{\mathbb{R}^{d-2}} [h((1-\varepsilon^2)^{1/2}v+\varepsilon y) - h(y)]Q(dv)\Phi_{d-2}(dy)| \\
= \quad & \sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} \chi_{\varepsilon^2}(v|h)Q(dv)|.
\end{aligned}
$$

The last equality uses (31). Finally using Lemma 11.4 of Bhattacharya and Rao (1986), page 95, we obtain

$$
\begin{aligned}
& \sup_{g\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} g(v)(Q-\Phi_{d-2})(dv)| \\
\le \quad & \sup_{g\in\mathcal{A}} \frac{1}{2\alpha-1}[\gamma^*(g;\varepsilon) + \alpha\tau^*(g;2\varepsilon) + (1-\alpha)\tau^*(g;\varepsilon)] \\
\le \quad & \frac{1}{2\alpha-1}\sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} h(v)(Q-\Phi_{d-2}) * \Phi_{d-2,\varepsilon/\sqrt{m_d(1-\varepsilon^2)}}(dv)| + \frac{(1+\alpha)\Delta\varepsilon}{2\alpha-1} \\
\le \quad & \frac{1}{(2\alpha-1)^2}\sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} h(v)(Q-\Phi_{d-2}) * \Phi_{d-2,\varepsilon\sqrt{2}/\sqrt{m_d(1-\varepsilon^2)}}(dv)| \\
& +(1+\alpha)\Delta\varepsilon[\frac{1}{2\alpha-1} + \frac{1}{(2\alpha-1)^2}] \\
\le \quad & \frac{1}{(2\alpha-1)^{m_d}}\sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} h(v)(Q-\Phi_{d-2}) * \Phi_{d-2,\varepsilon/\sqrt{1-\varepsilon^2}}(dv)| + \sum_{i=1}^{m_d}\frac{(1+\alpha)\Delta\varepsilon}{(2\alpha-1)^i} \\
= \quad & \frac{1}{(2\alpha-1)^{m_d}}\sup_{h\in\mathcal{A}} |\int_{\mathbb{R}^{d-2}} \chi_{\varepsilon^2}(v|h)Q(dv)| + \sum_{i=1}^{m_d}\frac{(1+\alpha)\Delta\varepsilon}{(2\alpha-1)^i}.
\end{aligned}
$$

This proves Lemma 2. □

The theorem below is the main result of this section and is needed in the proof of Theorem 2.

Theorem 3. *Suppose $d \geq 3$. Let $f : [0,1)^d \to \mathbb{R}$ be smooth with a Lipschitz continuous mixed partial of order d such that $\int_{\mathbb{R}^d} f_{rem}^2(x)dx > 0$ and the $(d-2)$-variate random vector V be as in (29). Then V converges to Φ_{d-2} in distribution as $q \to \infty$.*

Proof. In this proof it suffices to take $\Delta = 2\sqrt{d-2}$ in (30) and $\mathcal{A}$ to be the class of all indicator functions of measurable convex sets in $\mathbb{R}^{d-2}$. Let J be a random variable uniformly distributed over $\{1, \ldots, d\}$ and (B_1, B_2) be a random vector uniformly distributed over the set

$$\{(b_1, b_2) \in \{0, \ldots, q-1\}^2 : b_1 \neq b_2\}.$$

J and (B_1, B_2) are independent of each other and are also independent of all previously defined random quantities. Define for $j = 1, \ldots, d$,

$$\tilde{\pi}_j = \begin{cases} \pi_j & \text{if } j \neq J, \\ \tau_{B_1,B_2} \circ \pi_j & \text{if } j = J, \end{cases}$$

where τ_{B_1,B_2} denotes the permutation of $\{0, \ldots, q-1\}$ that transposes B_1 and B_2 leaving all other elements fixed. We further define for $\ell = 1, \ldots, d-2$,

$$\begin{aligned} \tilde{V}_\ell &= \frac{1}{q^2 \sigma_\ell} \sum_{i=1}^{q^2} \sum_{0 \leq u_1, \cdots, u_d \leq 1 : |u| = \ell+2} \nu^*[\tilde{\pi}_{j_1}(a_{i,j_1}); \ldots; \tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})], \\ \tilde{V} &= (\tilde{V}_1, \ldots, \tilde{V}_{d-2})'. \end{aligned} \tag{36}$$

From symmetry, we observe that $(V, \tilde{V})$ is an exchangeable pair of random vectors in that $(V, \tilde{V})$ and $(\tilde{V}, V)$ possess the same $2(d-2)$-variate distribution. We now write

$$\tilde{V}_\ell - V_\ell = \tilde{S}_\ell - S_\ell, \tag{37}$$

where

$$\begin{aligned} \tilde{S}_\ell = \frac{1}{q^2 \sigma_\ell} \sum_{i=1}^{q^2} \sum_{0 \leq u_1, \ldots, u_d \leq 1 : |u| = \ell+2} & \mathcal{I}\{J \in \{j_1, \ldots, j_{|u|}\}, \pi_J(a_{i,J}) \in \{B_1, B_2\}\} \\ & \times \nu^*[\tilde{\pi}_{j_1}(a_{i,j_1}); \ldots; \tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})], \end{aligned}$$

and

$$\begin{aligned} S_\ell = \frac{1}{q^2 \sigma_\ell} \sum_{i=1}^{q^2} \sum_{0 \leq u_1, \ldots, u_d \leq 1 : |u| = \ell+2} & \mathcal{I}\{J \in \{j_1, \ldots, j_{|u|}\}, \pi_J(a_{i,J}) \in \{B_1, B_2\}\} \\ & \times \nu^*[\pi_{j_1}(a_{i,j_1}); \ldots; \pi_{j_{|u|}}(a_{i,j_{|u|}})]. \end{aligned}$$

Let $\mathcal{W}$ be the σ-field generated by the random quantities

$$\{\pi_j(a_{i,j}) : i = 1, \ldots, q^2, j = 1, \ldots, d\},$$

$E^{\mathcal{W}}$ denote conditional expectation given $\mathcal{W}$ and $\psi_t(.)$ be as in (31). From the exchangeability of $(V, \tilde{V})$, we have for $0 < \varepsilon < 1/2$,

$$\begin{aligned} 0 &= E\{(\tilde{V}_i - V_i)[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V) + \frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(\tilde{V})]\} \\ &= 2E\{[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]E^{\mathcal{W}}(\tilde{V}_i - V_i)\} + E\{(\tilde{V}_i - V_i)[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(\tilde{V}) - \frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]\}. \end{aligned}$$

We observe from Proposition 3 (see Appendix) that

$$\begin{aligned} E\{(\tilde{V}_i - V_i)[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(\tilde{V}) - \frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]\} &= -2E\{[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]E^{\mathcal{W}}(\tilde{V}_i - V_i)\} \\ &= \frac{4(i+2)}{d(q-1)}E[V_i\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]. \end{aligned}$$

Now using Lemma 1, we have

$$\begin{aligned} & E[\chi_{\varepsilon^2}(V|h)] \\ = \;& E[\sum_{i=1}^{d-2} V_i\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V) - \sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] \\ = \;& \sum_{i=1}^{d-2}\frac{d(q-1)}{4(i+2)}E\{(\tilde{V}_i - V_i)[\frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(\tilde{V}) - \frac{\partial}{\partial v_i}\psi_{\varepsilon^2}(V)]\} - E[\sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] \\ = \;& \sum_{i=1}^{d-2}\sum_{j=1}^{d-2}\frac{d(q-1)}{4(i+2)}E\{(\tilde{V}_i - V_i)(\tilde{V}_j - V_j)\int_0^1\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V + t(\tilde{V} - V))dt\} \\ & -E[\sum_{i=1}^{d-2}\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] \\ = \;& \sum_{i=1}^{d-2}\Big\{\frac{d(q-1)}{4(i+2)}E\{(\tilde{V}_i - V_i)^2\int_0^1\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V + t(\tilde{V} - V))dt\} - E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]\Big\} \\ & +\sum_{i=1}^{d-2}\sum_{j\neq i}\frac{d(q-1)}{4(i+2)}E\{(\tilde{V}_i - V_i)(\tilde{V}_j - V_j)\int_0^1\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V + t(\tilde{V} - V))dt\} \\ = \;& \sum_{i=1}^{d-2}\Big\{[\frac{d(q-1)}{4(i+2)}E(\tilde{S}_i - S_i)^2 - 1]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] \\ & \quad -\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)^2]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] + \frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)^2\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)] \\ & \quad +\frac{d(q-1)}{4(i+2)}E\{(\tilde{S}_i - S_i)^2\int_0^1[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V + t(\tilde{V} - V)) - \frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]dt\}\Big\} \end{aligned}$$

$$
\begin{aligned}
&+\sum_{i=1}^{d-2}\sum_{j\neq i}\Big\{\frac{d(q-1)}{4(i+2)}E\{(\tilde{S}_i-S_i)(\tilde{S}_j-S_j)\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V)\}\\
&\quad+\frac{d(q-1)}{4(i+2)}E\{(\tilde{S}_i-S_i)(\tilde{S}_j-S_j)\\
&\quad\quad\times\int_0^1[\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V+t(\tilde{V}-V))-\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V)]dt\}\Big\}.
\end{aligned}
$$

Hence it follows from Propositions 4 to 7 (in the Appendix) that

$$E[\chi_{\varepsilon^2}(V|h)]=O(\frac{\log(1/\varepsilon)}{q^{1/2}})+O(\frac{1}{\varepsilon q^{1/2}}), \tag{38}$$

as $q\to\infty$ uniformly over $h\in\mathcal{A}$ and $\varepsilon\in(0,1/2)$. Using (38) and Lemma 2, we have

$$\sup_{g\in\mathcal{A}}|E[g(V)]-\int_{\mathbb{R}^{d-2}}g(v)\Phi_{d-2}(dv)|=O(\varepsilon+\frac{\log(1/\varepsilon)}{q^{1/2}}+\frac{1}{\varepsilon q^{1/2}}), \tag{39}$$

as $q\to\infty$ uniformly over $\varepsilon\in(0,1/2)$. By taking $\varepsilon=q^{-1/4}$, we conclude that the left hand side of (39) tends to 0 as $q\to\infty$. This implies that V converges to Φ_{d-2} in distribution as $q\to\infty$ and Theorem 3 is proved. □

4 Acknowledgements

I would like to thank Professor Rahul Mukerjee for his encouragement in the writing of this article. He has also spent quite a lot of time reading this article and spotting a number of oversights. For all these and more, I am very grateful to him.

5 Appendix

Lemma 3. *Let $f:[0,1)^d\to\mathbb{R}$ be smooth with a Lipschitz continuous mixed partial of order d. Then for $(a_1,\ldots,a_d)',(x_1,\ldots,x_d)'\in[0,1)^d$, we have*

$$
\begin{aligned}
f(x_1,\ldots,x_d) &= (x_1-a_1)\ldots(x_d-a_d)\frac{\partial^d f(a_1,\ldots,a_d)}{\partial x_1\ldots\partial x_d}\\
&\quad+\int_{a_1}^{x_1}\cdots\int_{a_d}^{x_d}[\frac{\partial^d f(t_1,\ldots,t_d)}{\partial x_1\ldots\partial x_d}-\frac{\partial^d f(a_1,\ldots,a_d)}{\partial x_1\ldots\partial x_d}]dt_d\ldots dt_1\\
&\quad+h_{1;a_1,\ldots,a_d}(x_2,\ldots,x_d)+h_{2;a_1,\ldots,a_d}(x_1,x_3,\ldots,x_d)+\cdots\\
&\quad+h_{d;a_1,\ldots,a_d}(x_1,\ldots,x_{d-1}),
\end{aligned} \tag{40}
$$

where $h_{i,a_1,\ldots,a_d}:\mathbb{R}^{d-1}\to\mathbb{R}$, $i=1,\ldots,d$ are suitably chosen functions.

PROOF. We shall use induction on d. Clearly, (40) holds for $d=1$ since in this case, we have

$$f(x_1)=f(a_1)+\int_{a_1}^{x_1}\frac{\partial f(t_1)}{\partial x_1}dt_1.$$

Now we assume that (40) holds for some $d=k\geq 1$. Then

$$\begin{aligned}
f(x_1,\ldots,x_{k+1}) &= \int_{a_1}^{x_1}\cdots\int_{a_k}^{x_k}\frac{\partial^k f(t_1,\ldots,t_k,x_{k+1})}{\partial x_1\ldots\partial x_k}dt_k\ldots dt_1\\
&\quad +h_{1;a_1,\ldots,a_k}(x_2,\ldots,x_k,x_{k+1})+h_{2;a_1,\ldots,a_k}(x_1,x_3,\ldots,x_k,x_{k+1})+\cdots\\
&\quad +h_{k;a_1,\ldots,a_k}(x_1,\ldots,x_{k-1},x_{k+1})\\
&= \int_{a_1}^{x_1}\cdots\int_{a_k}^{x_k}\frac{\partial^k f(t_1,\ldots,t_k,a_{k+1})}{\partial x_1\ldots\partial x_k}dt_k\ldots dt_1\\
&\qquad +\int_{a_1}^{x_1}\cdots\int_{a_{k+1}}^{x_{k+1}}\frac{\partial^{k+1} f(t_1,\ldots,t_{k+1})}{\partial x_1\ldots\partial x_{k+1}}dt_{k+1}\ldots dt_1\\
&\quad +h_{1;a_1,\ldots,a_k}(x_2,\ldots,x_k,x_{k+1})+h_{2;a_1,\ldots,a_k}(x_1,x_3,\ldots,x_k,x_{k+1})+\cdots\\
&\quad +h_{k;a_1,\ldots,a_k}(x_1,\ldots,x_{k-1},x_{k+1}).
\end{aligned}$$

This shows that (40) holds for $d=k+1$ and Lemma 3 is proved. □

Lemma 4. *Let $f:[0,1)^d\to\mathbb{R}$ be smooth with a Lipschitz continuous mixed partial of order d. Then*

$$\sum_{c_j=0}^{q-1}\langle f,\psi_{u_j-1,t_j,c_j}\rangle=0,\qquad \text{if } u_j\geq 1, \tag{41}$$

and

$$\begin{aligned}
\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle &= q^{|u|-\sum_{l=1}^{|u|}3u_{j_l}/2}\Big[\frac{\partial^{|u|}}{\partial x_{j_1}\ldots\partial x_{j_{|u|}}}f_{j_1,\ldots,j_{|u|}}\Big(\frac{t_{j_1}+0.5}{q^{u_{j_1}-1}},\ldots,\frac{t_{j_{|u|}}+0.5}{q^{u_{j_{|u|}}-1}}\Big)\Big]\\
&\quad\times\prod_{l=1}^{|u|}\Big(\frac{c_{j_l}}{q}+\frac{1}{2q}-\frac{1}{2}\Big)+O(1)q^{|u|-\sum_{l=1}^{|u|}3u_{j_l}/2}\max_{1\leq l\leq|u|}q^{-\beta(u_{j_l}-1)}
\end{aligned}\tag{42}$$

as $q\to\infty$ where

$$f_{j_1,\ldots,j_{|u|}}(x_{j_1},\ldots,x_{j_{|u|}})=\int_{[0,1)^{d-|u|}}f(x)\prod_{1\leq i\leq d:i\notin\{j_1,\ldots,j_{|u|}\}}dx_i.$$

Also,

$$\begin{aligned}
&E\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}),\pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\ldots,\pi_{i,j,u_j}(0):1\leq j\leq d]^2\}\\
&=\Big(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\Big)\cdots\Big(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\Big)\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2\sum_{k=0}^{|u|}\binom{|u|}{k}\frac{1}{q^k}\Big(1-\frac{1}{q}\Big)^{|u|-k},
\end{aligned}$$

and

$$
\begin{aligned}
(43)\qquad & E\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}),\pi_{i,j,2}(0),\ldots,\pi_{i,j,u_j}(0):1\le j\le d]^2\}\\
= \ & (\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2\sum_{k=0}^{|u|}\binom{|u|}{k}\frac{1}{q^k}(1-\frac{1}{q})^{|u|-k}.
\end{aligned}
$$

Finally $E\{\nu^*[\pi_{j_1}(a_{i,j_1});\ldots;\pi_{j_{|u|}}(a_{i,j_{|u|}})]^4\}=O(1)$ *as* $q\to\infty$.

PROOF. (41) easily follows from (11). To prove (42), we observe that

$$
\begin{aligned}
& \langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\\
= \ & \int_{[0,1)^d} f(x)[\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}(x_{j_l})]dx\\
= \ & \int_{[0,1)^d} f(x)\Big\{\prod_{l=1}^{|u|}q^{u_{j_l}/2}\Big[\mathcal{I}\{x_{j_l}\in[\frac{qt_{j_l}+c_{j_l}}{q^{u_{j_l}}},\frac{qt_{j_l}+c_{j_l}+1}{q^{u_{j_l}}})\}\\
& \quad -\frac{1}{q}\mathcal{I}\{x_{j_l}\in[\frac{t_{j_l}}{q^{u_{j_l}-1}},\frac{t_{j_l}+1}{q^{u_{j_l}-1}})\}\Big]\Big\}dx\\
= \ & q^{-\sum_{l=1}^{|u|}u_{j_l}/2}\int_{[0,1)^{|u|}} f_{j_1,\ldots,j_{|u|}}(x_{j_1},\ldots,x_{j_{|u|}})\Big\{\prod_{l=1}^{|u|}q^{u_{j_l}}\Big[\mathcal{I}\{x_{j_l}\in\\
(44)\qquad & \quad [\frac{qt_{j_l}+c_{j_l}}{q^{u_{j_l}}},\frac{qt_{j_l}+c_{j_l}+1}{q^{u_{j_l}}})\}-\frac{1}{q}\mathcal{I}\{x_{j_l}\in[\frac{t_{j_l}}{q^{u_{j_l}-1}},\frac{t_{j_l}+1}{q^{u_{j_l}-1}})\}\Big]\Big\}dx_{j_1}\ldots dx_{j_{|u|}}.
\end{aligned}
$$

We observe from Lemma 3 that the right hand side of (44) equals

$$
\begin{aligned}
& q^{-\sum_{l=1}^{|u|}u_{j_l}/2}[\frac{\partial^{|u|}}{\partial x_{j_1}\ldots\partial x_{j_{|u|}}}f_{j_1,\ldots,j_{|u|}}(\frac{t_{j_1}+0.5}{q^{u_{j_1}-1}},\ldots,\frac{t_{j_{|u|}}+0.5}{q^{u_{j_{|u|}}-1}})]\\
& \times\prod_{l=1}^{|u|}\Big[q^{u_{j_l}}\int_{(qt_{j_l}+c_{j_l})/q^{u_{j_l}}}^{(qt_{j_l}+c_{j_l}+1)/q^{u_{j_l}}}(x_{j_l}-\frac{t_{j_l}+0.5}{q^{u_{j_l}-1}})dx_{j_l}\\
& \quad -q^{u_{j_l}-1}\int_{t_{j_l}/q^{u_{j_l}-1}}^{(t_{j_l}+1)/q^{u_{j_l}-1}}(x_{j_l}-\frac{t_{j_l}+0.5}{q^{u_{j_l}-1}})dx_{j_l}\Big]\\
& +q^{-\sum_{l=1}^{|u|}u_{j_l}/2}\int_{[0,1)^{|u|}}\int_{(t_{j_1}+0.5)/q^{u_{j_1}-1}}^{x_{j_1}}\cdots\int_{(t_{j_{|u|}}+0.5)/q^{u_{j_{|u|}}-1}}^{x_{j_{|u|}}}\\
& \quad\times[\frac{\partial^{|u|}f_{j_1,\ldots,j_{|u|}}(s_1,\ldots,s_{|u|})}{\partial x_{j_1}\ldots\partial x_{j_{|u|}}}-\frac{\partial^{|u|}f_{j_1,\ldots,j_{|u|}}(\frac{t_{j_1}+0.5}{q^{u_{j_1}-1}},\ldots,\frac{t_{j_{|u|}}+0.5}{q^{u_{j_{|u|}}-1}})}{\partial x_{j_1}\ldots\partial x_{j_{|u|}}}]ds_{|u|}\ldots ds_1\\
& \quad\times\Big\{\prod_{l=1}^{|u|}q^{u_{j_l}}\Big[\mathcal{I}\{x_{j_l}\in[\frac{qt_{j_l}+c_{j_l}}{q^{u_{j_l}}},\frac{qt_{j_l}+c_{j_l}+1}{q^{u_{j_l}}})\}
\end{aligned}
$$

$$-\frac{1}{q}\mathcal{I}\{x_{j_l}\in[\frac{t_{j_l}}{q^{u_{j_l}-1}},\frac{t_{j_l}+1}{q^{u_{j_l}-1}})\}\Big]\Big\}dx_{j_1}\dots dx_{j_{|u|}}$$

$$= q^{|u|-\sum_{l=1}^{|u|}3u_{j_l}/2}[\frac{\partial^{|u|}}{\partial x_{j_1}\dots\partial x_{j_{|u|}}}f_{j_1,\dots,j_{|u|}}(\frac{t_{j_1}+0.5}{q^{u_{j_1}-1}},\dots,\frac{t_{j_{|u|}}+0.5}{q^{u_{j_{|u|}}-1}})]$$

$$\times\prod_{l=1}^{|u|}[\frac{c_{j_l}}{q}-\frac{1}{2}(1-\frac{1}{q})]+O(1)q^{|u|-\sum_{l=1}^{|u|}3u_{j_l}/2}\max_{1\le l\le|u|}q^{-\beta(u_{j_l}-1)},$$

as $q\to\infty$. Next we observe from (13) that

$$\nu_{u_1,\dots,u_d}[\pi_j(a_{i,j}),\pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\dots,\pi_{i,j,u_j}(0):1\le j\le d]$$

$$= (\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle$$

$$\times\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}+\frac{\pi_{j_l;a_{i,j_l}}(b_{i,j_l,2})}{q^2}+\sum_{k=3}^{u_{j_l}}\frac{\pi_{i,j_l,k}(0)}{q^k}),$$

and hence

$$E\{\nu_{u_1,\dots,u_d}[\pi_j(a_{i,j}),\pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\dots,\pi_{i,j,u_j}(0):1\le j\le d]^2\}$$

$$= (\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})(\sum_{t'_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c'_{j_1}=0}^{q-1})\cdots(\sum_{t'_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c'_{j_{|u|}}=0}^{q-1})$$

$$\times\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t'_{j_l},c'_{j_l}}\rangle$$

$$\times\prod_{l=1}^{|u|}E\Big[\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}+\frac{\pi_{j_l;a_{i,j_l}}(b_{i,j_l,2})}{q^2}+\sum_{k=3}^{u_{j_l}}\frac{\pi_{i,j_l,k}(0)}{q^k})$$

$$\times\psi_{u_{j_l}-1,t'_{j_l},c'_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}+\frac{\pi_{j_l;a_{i,j_l}}(b_{i,j_l,2})}{q^2}+\sum_{k=3}^{u_{j_l}}\frac{\pi_{i,j_l,k}(0)}{q^k})\Big]$$

$$= (\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\sum_{c'_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\sum_{c'_{j_{|u|}}=0}^{q-1})$$

$$\times\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c'_{j_l}}\rangle$$

$$\times\prod_{l=1}^{|u|}E\Big[\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}+\frac{\pi_{j_l;a_{i,j_l}}(b_{i,j_l,2})}{q^2}+\sum_{k=3}^{u_{j_l}}\frac{\pi_{i,j_l,k}(0)}{q^k})$$

$$\times\psi_{u_{j_l}-1,t_{j_l},c'_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}+\frac{\pi_{j_l;a_{i,j_l}}(b_{i,j_l,2})}{q^2}+\sum_{k=3}^{u_{j_l}}\frac{\pi_{i,j_l,k}(0)}{q^k})\Big].$$

We observe that the right hand side of the last equation can be expressed as a finite sum of terms of the following form (up to permutations): for $\alpha = 0, \ldots, |u|$,

$$
\begin{aligned}
&\Big(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\sum_{c'_{j_1}\neq c_{j_1}}\Big)\cdots\Big(\sum_{t_{j_\alpha}=0}^{q^{u_{j_\alpha}-1}-1}\sum_{c_{j_\alpha}=0}^{q-1}\sum_{c'_{j_\alpha}\neq c_{j_\alpha}}\Big)\\
&\times\Big(\sum_{t_{j_{\alpha+1}}=0}^{q^{u_{j_{\alpha+1}}-1}-1}\sum_{c_{j_{\alpha+1}}=0}^{q-1}\Big)\cdots\Big(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\Big)\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\\
&\times\langle f,\Big(\prod_{l_1=1}^{\alpha}\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c'_{j_{l_1}}}\Big)\Big(\prod_{l_2=\alpha+1}^{|u|}\psi_{u_{j_{l_2}}-1,t_{j_{l_2}},c_{j_{l_2}}}\Big)\rangle\\
&\times E\Big[\prod_{l_1=1}^{\alpha}\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c_{j_{l_1}}}\Big(\frac{\pi_{j_{l_1}}(a_{i,j_{l_1}})}{q}+\frac{\pi_{j_{l_1};a_{i,j_{l_1}}}(b_{i,j_{l_1},2})}{q^2}+\sum_{k=3}^{u_{j_{l_1}}}\frac{\pi_{i,j_{l_1},k}(0)}{q^k}\Big)\\
&\quad\times\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c'_{j_{l_1}}}\Big(\frac{\pi_{j_{l_1}}(a_{i,j_{l_1}})}{q}+\frac{\pi_{j_{l_1};a_{i,j_{l_1}}}(b_{i,j_{l_1},2})}{q^2}+\sum_{k=3}^{u_{j_{l_1}}}\frac{\pi_{i,j_{l_1},k}(0)}{q^k}\Big)\Big]\\
&\times E\Big[\prod_{l_2=\alpha+1}^{|u|}\psi_{u_{j_{l_2}}-1,t_{j_{l_2}},c_{j_{l_2}}}\Big(\frac{\pi_{j_{l_2}}(a_{i,j_{l_2}})}{q}+\frac{\pi_{j_{l_2};a_{i,j_{l_2}}}(b_{i,j_{l_2},2})}{q^2}+\sum_{k=3}^{u_{j_{l_2}}}\frac{\pi_{i,j_{l_2},k}(0)}{q^k}\Big)^2\Big]\\
=\;&\Big(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\sum_{c'_{j_1}\neq c_{j_1}}\Big)\cdots\Big(\sum_{t_{j_\alpha}=0}^{q^{u_{j_\alpha}-1}-1}\sum_{c_{j_\alpha}=0}^{q-1}\sum_{c'_{j_\alpha}\neq c_{j_\alpha}}\Big)\\
&\quad\times\Big(\sum_{t_{j_{\alpha+1}}=0}^{q^{u_{j_{\alpha+1}}-1}-1}\sum_{c_{j_{\alpha+1}}=0}^{q-1}\Big)\cdots\Big(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1}\Big)\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\\
&\quad\times\langle f,\Big(\prod_{l_1=1}^{\alpha}\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c'_{j_{l_1}}}\Big)\Big(\prod_{l_2=\alpha+1}^{|u|}\psi_{u_{j_{l_2}}-1,t_{j_{l_2}},c_{j_{l_2}}}\Big)\rangle E\Big[\prod_{l_1=1}^{\alpha}\Big(\\
&\qquad -q^{u_{j_{l_1}}-1}\mathcal{I}\{\frac{qt_{j_{l_1}}+c_{j_{l_1}}}{q^{u_{j_{l_1}}}}\le\frac{\pi_{j_{l_1}}(a_{i,j_{l_1}})}{q}+\frac{\pi_{j_{l_1};a_{i,j_{l_1}}}(b_{i,j_{l_1},2})}{q^2}\\
&\qquad\quad+\sum_{k=3}^{u_{j_{l_1}}}\frac{\pi_{i,j_{l_1},k}(0)}{q^k}<\frac{qt_{j_{l_1}}+c_{j_{l_1}}+1}{q^{u_{j_{l_1}}}}\}\\
&\qquad -q^{u_{j_{l_1}}-1}\mathcal{I}\{\frac{qt_{j_{l_1}}+c'_{j_{l_1}}}{q^{u_{j_{l_1}}}}\le\frac{\pi_{j_{l_1}}(a_{i,j_{l_1}})}{q}+\frac{\pi_{j_{l_1};a_{i,j_{l_1}}}(b_{i,j_{l_1},2})}{q^2}\\
&\qquad\quad+\sum_{k=3}^{u_{j_{l_1}}}\frac{\pi_{i,j_{l_1},k}(0)}{q^k}<\frac{qt_{j_{l_1}}+c'_{j_{l_1}}+1}{q^{u_{j_{l_1}}}}\}\\
&\qquad +q^{u_{j_{l_1}}-2}\mathcal{I}\{\frac{t_{j_{l_1}}}{q^{u_{j_{l_1}}-1}}\le\frac{\pi_{j_{l_1}}(a_{i,j_{l_1}})}{q}+\frac{\pi_{j_{l_1};a_{i,j_{l_1}}}(b_{i,j_{l_1},2})}{q^2}
\end{aligned}
$$

$$
\begin{aligned}
&\qquad +\sum_{k=3}^{u_{j_{l_1}}}\frac{\pi_{i,j_{l_1},k}(0)}{q^k}<\frac{t_{j_{l_1}}+1}{q^{u_{j_{l_1}}-1}}\}\Big)\Big]\\
&\quad\times E\Big[\prod_{l_2=\alpha+1}^{|u|}\Big(q^{u_{j_{l_2}}}\mathcal{I}\{\frac{qt_{j_{l_2}}+c_{j_{l_2}}}{q^{u_{j_{l_2}}}}\le\frac{\pi_{j_{l_2}}(a_{i,j_{l_2}})}{q}+\frac{\pi_{j_{l_2};a_{i,j_{l_2}}}(b_{i,j_{l_2},2})}{q^2}\\
&\qquad +\sum_{k=3}^{u_{j_{l_2}}}\frac{\pi_{i,j_{l_2},k}(0)}{q^k}<\frac{qt_{j_{l_2}}+c_{j_{l_2}}+1}{q^{u_{j_{l_2}}}}\}\\
&\quad -2q^{u_{j_{l_2}}-1}\mathcal{I}\{\frac{qt_{j_{l_2}}+c_{j_{l_2}}}{q^{u_{j_{l_2}}}}\le\frac{\pi_{j_{l_2}}(a_{i,j_{l_2}})}{q}+\frac{\pi_{j_{l_2};a_{i,j_{l_2}}}(b_{i,j_{l_2},2})}{q^2}\\
&\qquad +\sum_{k=3}^{u_{j_{l_2}}}\frac{\pi_{i,j_{l_2},k}(0)}{q^k}<\frac{qt_{j_{l_2}}+c_{j_{l_2}}+1}{q^{u_{j_{l_2}}}}\}\\
&\quad +q^{u_{j_{l_2}}-2}\mathcal{I}\{\frac{t_{j_{l_2}}}{q^{u_{j_{l_2}}-1}}\le\frac{\pi_{j_{l_2}}(a_{i,j_{l_2}})}{q}+\frac{\pi_{j_{l_2};a_{i,j_{l_2}}}(b_{i,j_{l_2},2})}{q^2}\\
&\qquad +\sum_{k=3}^{u_{j_{l_2}}}\frac{\pi_{i,j_{l_2},k}(0)}{q^k}<\frac{t_{j_{l_2}}+1}{q^{u_{j_{l_2}}-1}}\}\Big)\Big]\\
={}&(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\sum_{c'_{j_1}\ne c_{j_1}})\cdots(\sum_{t_{j_\alpha}=0}^{q^{u_{j_\alpha}-1}-1}\sum_{c_{j_\alpha}=0}^{q-1}\sum_{c'_{j_\alpha}\ne c_{j_\alpha}})\\
&\times(\sum_{t_{j_{\alpha+1}}=0}^{q^{u_{j_{\alpha+1}}-1}-1}\sum_{c_{j_{\alpha+1}}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\\
&\times\langle f,(\prod_{l_1=1}^{\alpha}\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c'_{j_{l_1}}})(\prod_{l_2=\alpha+1}^{|u|}\psi_{u_{j_{l_2}}-1,t_{j_{l_2}},c_{j_{l_2}}})\rangle(-\frac{1}{q})^{\alpha}(1-\frac{1}{q})^{|u|-\alpha}\\
={}&(-\frac{1}{q})^{\alpha}(1-\frac{1}{q})^{|u|-\alpha}[\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1}\\
&\times(\sum_{c'_{j_1}=0}^{q-1}-\mathcal{I}\{c'_{j_1}=c_{j_1}\})]\cdots[\sum_{t_{j_\alpha}=0}^{q^{u_{j_\alpha}-1}-1}\sum_{c_{j_\alpha}=0}^{q-1}(\sum_{c'_{j_\alpha}=0}^{q-1}-\mathcal{I}\{c'_{j_\alpha}=c_{j_\alpha}\})]\\
&\times(\sum_{t_{j_{\alpha+1}}=0}^{q^{u_{j_{\alpha+1}}-1}-1}\sum_{c_{j_{\alpha+1}}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle\\
&\times\langle f,(\prod_{l_1=1}^{\alpha}\psi_{u_{j_{l_1}}-1,t_{j_{l_1}},c'_{j_{l_1}}})(\prod_{l_2=\alpha+1}^{|u|}\psi_{u_{j_{l_2}}-1,t_{j_{l_2}},c_{j_{l_2}}})\rangle
\end{aligned}
$$

$$= \frac{1}{q^{\alpha}}(1-\frac{1}{q})^{|u|-\alpha}(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2.$$

Finally we conclude via symmetry that

$$\begin{aligned}
& E\{\nu_{u_1,\ldots,u_d}[\pi_j(a_{i,j}),\pi_{j;a_{i,j}}(b_{i,j,2}),\pi_{i,j,3}(0),\ldots,\pi_{i,j,u_j}(0):1\le j\le d]^2\} \\
= & \sum_{\alpha=0}^{|u|}\binom{|u|}{\alpha}\frac{1}{q^{\alpha}}(1-\frac{1}{q})^{|u|-\alpha}(\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}}\rangle^2.
\end{aligned}$$

The proof of (43) is similar and is omitted. Finally using (13), (17) and that $u_{j_1}=\cdots=u_{j_{|u|}}=1$, we have

$$\begin{aligned}
& E\{\nu^*[\pi_{j_1}(a_{i,j_1});\ldots;\pi_{j_{|u|}}(a_{i,j_{|u|}})]^4\} \\
= & (\sum_{t_{j_1}=0}^{q^{u_{j_1}-1}-1}\sum_{c_{j_1}^{(1)}=0}^{q-1}\sum_{c_{j_1}^{(2)}=0}^{q-1}\sum_{c_{j_1}^{(3)}=0}^{q-1}\sum_{c_{j_1}^{(4)}=0}^{q-1})\cdots(\sum_{t_{j_{|u|}}=0}^{q^{u_{j_{|u|}}-1}-1}\sum_{c_{j_{|u|}}^{(1)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(2)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(3)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(4)}=0}^{q-1}) \\
& \times\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(1)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(2)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(3)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(4)}}\rangle \\
& \times\prod_{l=1}^{|u|}E\Big[\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(2)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}) \\
& \quad\times\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(3)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{u_{j_l}-1,t_{j_l},c_{j_l}^{(4)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big] \\
= & (\sum_{c_{j_1}^{(1)}=0}^{q-1}\sum_{c_{j_1}^{(2)}=0}^{q-1}\sum_{c_{j_1}^{(3)}=0}^{q-1}\sum_{c_{j_1}^{(4)}=0}^{q-1})\cdots(\sum_{c_{j_{|u|}}^{(1)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(2)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(3)}=0}^{q-1}\sum_{c_{j_{|u|}}^{(4)}=0}^{q-1})\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(1)}}\rangle \\
& \times\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(2)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(3)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(4)}}\rangle\prod_{l=1}^{|u|}E[\prod_{k=1}^{4}\psi_{0,0,c_{j_l}^{(k)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})]. \qquad (45)
\end{aligned}$$

It is convenient to define the following subsets of $\{1,\ldots,d\}$: for $\{l_1,l_2,l_3,l_4\}=\{1,2,3,4\}$,

$$\begin{aligned}
\Xi_{\{1,2,3,4\}} &= \{l\in\{1,\ldots,|u|\}:c_{j_l}^{(1)}=c_{j_l}^{(2)}=c_{j_l}^{(3)}=c_{j_l}^{(4)}\}, \\
\Xi_{\{l_1,l_2,l_3\},\{l_4\}} &= \{l\in\{1,\ldots,|u|\}:c_{j_l}^{(l_1)}=c_{j_l}^{(l_2)}=c_{j_l}^{(l_3)}\neq c_{j_l}^{(l_4)}\}, \\
\Xi_{\{l_1,l_2\},\{l_3\},\{l_4\}} &= \{l\in\{1,\ldots,|u|\}:c_{j_l}^{(l_1)}=c_{j_l}^{(l_2)},\text{and } c_{j_l}^{(l_1)},c_{j_l}^{(l_3)},c_{j_l}^{(l_4)}\text{ are all distinct}\}, \\
\Xi_{\{l_1,l_2\},\{l_3,l_4\}} &= \{l\in\{1,\ldots,|u|\}:c_{j_l}^{(l_1)}=c_{j_l}^{(l_2)}\neq c_{j_l}^{(l_3)}=c_{j_l}^{(l_4)}\}, \\
\Xi_{\{l_1\},\{l_2\},\{l_3\},\{l_4\}} &= \{l\in\{1,\ldots,|u|\}:c_{j_l}^{(l_1)},c_{j_l}^{(l_2)},c_{j_l}^{(l_3)},c_{j_l}^{(l_4)}\text{ are all distinct}\}.
\end{aligned}$$

The right hand side of (45) can be written as a finite sum of terms of the following form:

$$
\begin{aligned}
&[\prod_{l\in\Xi_{\{1,2,3,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}][\prod_{l\in\Xi_{\{1,2,3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\neq c_{j_l}^{(1)}}]\\
&\times[\prod_{l\in\Xi_{\{1,2,4\},\{3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1,3,4\},\{2\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}]\\
&\times[\prod_{l\in\Xi_{\{2,3,4\},\{1\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\neq c_{j_l}^{(2)}}]\\
&\times[\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\neq c_{j_l}^{(1)},c_{j_l}^{(3)}}]\\
&\times[\prod_{l\in\Xi_{\{1,3\},\{2\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\neq c_{j_l}^{(1)},c_{j_l}^{(2)}}]\\
&\times[\prod_{l\in\Xi_{\{1,4\},\{2\},\{3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)},c_{j_l}^{(3)}}]\\
&\times[\prod_{l\in\Xi_{\{2,3\},\{1\},\{4\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\neq c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\neq c_{j_l}^{(1)},c_{j_l}^{(2)}}]\\
&\times[\prod_{l\in\Xi_{\{2,4\},\{1\},\{3\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\neq c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)},c_{j_l}^{(2)}}]\\
&\times[\prod_{l\in\Xi_{\{3,4\},\{1\},\{2\}}}\sum_{c_{j_l}^{(3)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\neq c_{j_l}^{(3)}}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)},c_{j_l}^{(3)}}]\\
&\times[\prod_{l\in\Xi_{\{1,2\},\{3,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1,3\},\{2,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}]\\
&\times[\prod_{l\in\Xi_{\{1,4\},\{2,3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\neq c_{j_l}^{(1)}}\\
&\quad\times\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\neq c_{j_l}^{(1)},c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\neq c_{j_l}^{(1)},c_{j_l}^{(2)},c_{j_l}^{(3)}}]\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(1)}}\rangle
\end{aligned}
$$

$$
\begin{aligned}
&\times\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c^{(2)}_{j_l}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c^{(3)}_{j_l}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c^{(4)}_{j_l}}\rangle\Big\{\prod_{l\in\Xi_{\{1,2,3,4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^4\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,2,3\},\{4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^3\psi_{0,0,c^{(4)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,2,4\},\{3\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^3\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,3,4\},\{2\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^3\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{2,3,4\},\{1\}}}E\Big[\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^3\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(4)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,3\},\{2\},\{4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(4)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,4\},\{2\},\{3\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{2,3\},\{1\},\{4\}}}E\Big[\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(4)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{2,4\},\{1\},\{3\}}}E\Big[\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{3,4\},\{1\},\{2\}}}E\Big[\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,2\},\{3,4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,3\},\{2,4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1,4\},\{2,3\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\Big]\Big\}\\
&\times\Big\{\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}}E\Big[\psi_{0,0,c^{(1)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(2)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\\
&\qquad\times\psi_{0,0,c^{(3)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c^{(4)}_{j_l}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\Big\}.
\end{aligned}
\tag{46}
$$

Now,

$$\begin{aligned}
&\prod_{l\in\Xi_{\{1,2,3,4\}}} E\Big[\psi_{0,0,c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^4\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3,4\}}} E\Big[[q^{1/2}\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}-q^{-1/2}]^4\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3,4\}}} q^2E\Big[[\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}-\frac{1}{q}]^4\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3,4\}}} q^2E\Big[(1-\frac{4}{q}+\frac{6}{q^2}-\frac{4}{q^3})\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}+\frac{1}{q^4}\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3,4\}}} (q-4+\frac{6}{q}-\frac{3}{q^2}),
\end{aligned}$$

$$\begin{aligned}
&\prod_{l\in\Xi_{\{1,2,3\},\{4\}}} E\Big[\psi_{0,0,c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^3\psi_{0,0,c_{j_l}^{(4)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3\},\{4\}}} q^2E\Big[[\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}-\frac{1}{q}]^3\\
&\times[\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(4)}+1}{q}\}-\frac{1}{q}]\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3\},\{4\}}} q^2E\Big[[(1-\frac{3}{q}+\frac{3}{q^2})\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}-\frac{1}{q^3}]\\
&\times[\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(4)}+1}{q}\}-\frac{1}{q}]\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3\},\{4\}}} q^2E\Big[-(\frac{1}{q}-\frac{3}{q^2}+\frac{3}{q^3})\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}+\frac{1}{q^4}\\
&-\frac{1}{q^3}\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(4)}+1}{q}\}\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2,3\},\{4\}}} (-1+\frac{3}{q}-\frac{3}{q^2}),
\end{aligned}$$

$$\begin{aligned}
&\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}} E\Big[\psi_{0,0,c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c_{j_l}^{(3)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c_{j_l}^{(4)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big]\\
=\ &\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}} q^2E\Big[[\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q}\le\frac{\pi_{j_l}(a_{i,j_l})}{q}<\frac{c_{j_l}^{(1)}+1}{q}\}-\frac{1}{q}]^2
\end{aligned}$$

$$
\begin{aligned}
&\quad \times[\mathcal{I}\{\frac{c_{j_l}^{(3)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(3)}+1}{q}\} - \frac{1}{q}][\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(4)}+1}{q}\} - \frac{1}{q}]\Big] \\
= &\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}} E\Big[(1-\frac{2}{q})\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(1)}+1}{q}\} + \frac{1}{q^2} \\
&\quad -\frac{1}{q}\mathcal{I}\{\frac{c_{j_l}^{(3)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(3)}+1}{q}\} - \frac{1}{q}\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(4)}+1}{q}\}\Big] \\
= &\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}} (\frac{1}{q} - \frac{3}{q^2}), \\
&\prod_{l\in\Xi_{\{1,2\},\{3,4\}}} E\Big[\psi_{0,0,c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\psi_{0,0,c_{j_l}^{(3)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})^2\Big]\Big\} \\
= &\prod_{l\in\Xi_{\{1,2\},\{3,4\}}} q^2E\Big[[\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(1)}+1}{q}\} - \frac{1}{q}]^2 \\
&\quad \times[\mathcal{I}\{\frac{c_{j_l}^{(3)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(3)}+1}{q}\} - \frac{1}{q}]^2\Big] \\
= &\prod_{l\in\Xi_{\{1,2\},\{3,4\}}} E\Big[(1-\frac{2}{q})\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(1)}+1}{q}\} \\
&\quad +(1-\frac{2}{q})\mathcal{I}\{\frac{c_{j_l}^{(3)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(3)}+1}{q}\} + \frac{1}{q^2}\Big] \\
= &\prod_{l\in\Xi_{\{1,2\},\{3,4\}}} (\frac{2}{q} - \frac{3}{q^2}), \\
&\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}} E\Big[\psi_{0,0,c_{j_l}^{(1)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c_{j_l}^{(2)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q}) \\
&\quad \times\psi_{0,0,c_{j_l}^{(3)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\psi_{0,0,c_{j_l}^{(4)}}(\frac{\pi_{j_l}(a_{i,j_l})}{q})\Big] \\
= &\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}} q^2E\Big[[\mathcal{I}\{\frac{c_{j_l}^{(1)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(1)}+1}{q}\} - \frac{1}{q}] \\
&\quad \times[\mathcal{I}\{\frac{c_{j_l}^{(2)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(2)}+1}{q}\} - \frac{1}{q}][\mathcal{I}\{\frac{c_{j_l}^{(3)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(3)}+1}{q}\} - \frac{1}{q}] \\
&\quad \times[\mathcal{I}\{\frac{c_{j_l}^{(4)}}{q} \le \frac{\pi_{j_l}(a_{i,j_l})}{q} < \frac{c_{j_l}^{(4)}+1}{q}\} - \frac{1}{q}]\Big] \\
= &\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}} (-\frac{3}{q^2}).
\end{aligned}
$$

Writing $|A|$ to denote the cardinality of a finite set A, we observe that (46) is equal to

$$[\prod_{l\in\Xi_{\{1,2,3,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}][\prod_{l\in\Xi_{\{1,2,3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\ne c_{j_l}^{(1)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,2,4\},\{3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1,3,4\},\{2\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}]$$

$$\times[\prod_{l\in\Xi_{\{2,3,4\},\{1\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\ne c_{j_l}^{(2)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,2\},\{3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\ne c_{j_l}^{(1)},c_{j_l}^{(3)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,3\},\{2\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\ne c_{j_l}^{(1)},c_{j_l}^{(2)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,4\},\{2\},\{3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)},c_{j_l}^{(3)}}]$$

$$\times[\prod_{l\in\Xi_{\{2,3\},\{1\},\{4\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\ne c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\ne c_{j_l}^{(1)},c_{j_l}^{(2)}}]$$

$$\times[\prod_{l\in\Xi_{\{2,4\},\{1\},\{3\}}}\sum_{c_{j_l}^{(2)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\ne c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)},c_{j_l}^{(2)}}]$$

$$\times[\prod_{l\in\Xi_{\{3,4\},\{1\},\{2\}}}\sum_{c_{j_l}^{(3)}=0}^{q-1}\sum_{0\le c_{j_l}^{(1)}\le q-1:c_{j_l}^{(1)}\ne c_{j_l}^{(3)}}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)},c_{j_l}^{(3)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,2\},\{3,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1,3\},\{2,4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}]$$

$$\times[\prod_{l\in\Xi_{\{1,4\},\{2,3\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}][\prod_{l\in\Xi_{\{1\},\{2\},\{3\},\{4\}}}\sum_{c_{j_l}^{(1)}=0}^{q-1}\sum_{0\le c_{j_l}^{(2)}\le q-1:c_{j_l}^{(2)}\ne c_{j_l}^{(1)}}$$

$$\times\sum_{0\le c_{j_l}^{(3)}\le q-1:c_{j_l}^{(3)}\ne c_{j_l}^{(1)},c_{j_l}^{(2)}}\sum_{0\le c_{j_l}^{(4)}\le q-1:c_{j_l}^{(4)}\ne c_{j_l}^{(1)},c_{j_l}^{(2)},c_{j_l}^{(3)}}]\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(1)}}\rangle$$

$$\begin{aligned}
&\times\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(2)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(3)}}\rangle\langle f,\prod_{l=1}^{|u|}\psi_{0,0,c_{j_l}^{(4)}}\rangle(q-4+\frac{6}{q}-\frac{3}{q^2})^{|\Xi_{\{1,2,3,4\}}|}\\
&\times(-1+\frac{3}{q}-\frac{3}{q^2})^{|\Xi_{\{1,2,3\},\{4\}}|+|\Xi_{\{1,2,4\},\{3\}}|+\Xi_{\{1,3,4\},\{2\}}|+\Xi_{\{2,3,4\},\{1\}}|}\\
&\times(\frac{1}{q}-\frac{3}{q^2})^{|\Xi_{\{1,2\},\{3\},\{4\}}|+|\Xi_{\{1,3\},\{2\},\{4\}}|+\Xi_{\{1,4\},\{2\},\{3\}}|+|\Xi_{\{2,3\},\{1\},\{4\}}|+|\Xi_{\{2,4\},\{1\},\{3\}}|+|\Xi_{\{3,4\},\{1\},\{2\}}|}\\
&\times(\frac{2}{q}-\frac{3}{q^2})^{|\Xi_{\{1,2\},\{3,4\}}|+|\Xi_{\{1,3\},\{2,4\}}|+|\Xi_{\{1,4\},\{2,3\}}|}(-\frac{3}{q^2})^{|\Xi_{\{1\},\{2\},\{3\},\{4\}}|}\\
= {}& O(1)q^{|\Xi_{\{1,2,3,4\}}|}q^{2(|\Xi_{\{1,2,3\},\{4\}}|+|\Xi_{\{1,2,4\},\{3\}}|+|\Xi_{\{1,3,4\},\{2\}}|+|\Xi_{\{2,3,4\},\{1\}}|)}\\
&\times q^{3(|\Xi_{\{1,2\},\{3\},\{4\}}|+|\Xi_{\{1,3\},\{2\},\{4\}}|+|\Xi_{\{1,4\},\{2\},\{3\}}|+|\Xi_{\{2,3\},\{1\},\{4\}}|+|\Xi_{\{2,4\},\{1\},\{3\}}|+|\Xi_{\{3,4\},\{1\},\{2\}}|)}\\
&\times q^{2(|\Xi_{\{1,2\},\{3,4\}}|+\Xi_{\{1,3\},\{2,4\}}|+|\Xi_{\{1,4\},\{2,3\}}|)}q^{4|\Xi_{\{1\},\{2\},\{3\},\{4\}}|}q^{-2|u|}q^{|\Xi_{\{1,2,3,4\}}|}\\
&\times(\frac{1}{q})^{|\Xi_{\{1,2\},\{3\},\{4\}}|+|\Xi_{\{1,3\},\{2\},\{4\}}|+|\Xi_{\{1,4\},\{2\},\{3\}}|+|\Xi_{\{2,3\},\{1\},\{4\}}|+|\Xi_{\{2,4\},\{1\},\{3\}}|+|\Xi_{\{3,4\},\{1\},\{2\}}|)}\\
&\times(\frac{2}{q})^{|\Xi_{\{1,2\},\{3,4\}}|+\Xi_{\{1,3\},\{2,4\}}|+|\Xi_{\{1,4\},\{2,3\}}}(\frac{3}{q^2})^{|\Xi_{\{1\},\{2\},\{3\},\{4\}}|}\\
= {}& O(1),
\end{aligned}$$

as $q \to \infty$. The last equality uses (42). This proves Lemma 4. □

Proposition 3. *Let V_ℓ and $\tilde{V}_\ell, \ell = 1, \ldots, d-2$, be as in (29) and (36) respectively. Then*

$$E^{\mathcal{W}}(\tilde{V}_\ell - V_\ell) = -\frac{2(\ell+2)}{d(q-1)}V_\ell.$$

PROOF. First we observe from (37) that

$$\begin{aligned}
E^{\mathcal{W}}(S_\ell) &= E^{\mathcal{W}}\Big\{\frac{1}{dq^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{k \in \{j_1,\ldots,j_{|u|}\}\}\\
&\qquad\times\mathcal{I}\{\pi_k(a_{i,k}) \in \{B_1,B_2\}\}\nu^*[\pi_{j_1}(a_{i,j_1});\cdots;\pi_{j_{|u|}}(a_{i,j_{|u|}})]\Big\}\\
&= \frac{2(\ell+2)}{dq^3\sigma_\ell}\sum_{i=1}^{q^2}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\nu^*[\pi_{j_1}(a_{i,j_1});\cdots;\pi_{j_{|u|}}(a_{i,j_{|u|}})],
\end{aligned}$$

and

$$\begin{aligned}
E^{\mathcal{W}}(\tilde{S}_\ell) &= E^{\mathcal{W}}\Big\{\frac{1}{dq^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{k \in \{j_1,\ldots,j_{|u|}\}\}\\
&\qquad\times\mathcal{I}\{\pi_k(a_{i,k}) \in \{B_1,B_2\}\}\nu^*[\tilde{\pi}_{j_1}(a_{i,j_1});\cdots;\tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})]\Big\}\\
&= E^{\mathcal{W}}\Big\{\frac{1}{dq^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\sum_{l=1}^{|u|}\mathcal{I}\{k = j_l\}
\end{aligned}$$

$$
\begin{aligned}
& \times\mathcal{I}\{\pi_{j_l}(a_{i,j_l}) \in \{B_1, B_2\}\}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{l-1}}(a_{i,j_{l-1}}); \\
& \qquad \tau_{B_1,B_2} \circ \pi_{j_l}(a_{i,j_l}); \pi_{j_{l+1}}(a_{i,j_{l+1}}); \cdots ; \tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})]\Big\} \\
= {} & E^{\mathcal{W}}\Big\{\frac{1}{dq^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\sum_{l=1}^{|u|}\mathcal{I}\{k=j_l\} \\
& \quad \times\mathcal{I}\{\pi_{j_l}(a_{i,j_l}) = B_1\}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{l-1}}(a_{i,j_{l-1}}); \\
& \qquad B_2; \pi_{j_{l+1}}(a_{i,j_{l+1}}); \cdots ; \tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})]\Big\} \\
& +E^{\mathcal{W}}\Big\{\frac{1}{dq^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\sum_{l=1}^{|u|}\mathcal{I}\{k=j_l\} \\
& \quad \times\mathcal{I}\{\pi_{j_l}(a_{i,j_l}) = B_2\}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{l-1}}(a_{i,j_{l-1}}); \\
& \qquad B_1; \pi_{j_{l+1}}(a_{i,j_{l+1}}); \cdots ; \tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})]\Big\} \\
= {} & -E^{\mathcal{W}}\Big\{\frac{1}{dq^2(q-1)\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\sum_{l=1}^{|u|}\mathcal{I}\{k=j_l\} \\
& \quad \times\mathcal{I}\{\pi_{j_l}(a_{i,j_l}) = B_1\}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{|u|}}(a_{i,j_{|u|}})]\Big\} \\
& -E^{\mathcal{W}}\Big\{\frac{1}{dq^2(q-1)\sigma_\ell}\sum_{i=1}^{q^2}\sum_{k=1}^{d}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\sum_{l=1}^{|u|}\mathcal{I}\{k=j_l\} \\
& \quad \times\mathcal{I}\{\pi_{j_l}(a_{i,j_l}) = B_2\}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{|u|}}(a_{i,j_{|u|}})]\Big\} \\
= {} & -\frac{2(\ell+2)}{dq^3(q-1)\sigma_\ell}\sum_{i=1}^{q^2}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{|u|}}(a_{i,j_{|u|}})].
\end{aligned}
$$

Thus we conclude that

$$
\begin{aligned}
E^{\mathcal{W}}(\tilde{V}_\ell - V_\ell) &= E^{\mathcal{W}}(\tilde{S}_\ell - S_\ell) \\
&= -\frac{2(\ell+2)}{dq^2(q-1)\sigma_\ell}\sum_{i=1}^{q^2}\sum_{0\le u_1,\cdots,u_d\le 1:|u|=\ell+2}\nu^*[\pi_{j_1}(a_{i,j_1}); \ldots ; \pi_{j_{|u|}}(a_{i,j_{|u|}})] \\
&= -\frac{2(\ell+2)}{d(q-1)}V_\ell. \qquad \square
\end{aligned}
$$

Proposition 4. *Let* S_i *and* $\tilde{S}_i$, $i = 1, \ldots, d-2$, *be as in (37). Then*

$$|\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)^2] - 1| = O(1/q), \qquad \forall i = 1, \ldots, d-2,$$

and

$$|\sum_{i=1}^{d-2}\{[\frac{d(q-1)}{4(i+2)}E(\tilde{S}_i - S_i)^2 - 1]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]\}| \le O(\frac{\|h\|_\infty}{q})\log(1/\varepsilon), \tag{47}$$

as $q \to \infty$ *uniformly over* $0 < \varepsilon < 1$.

Proof. For $\ell = 1, \dots, d-2$ and $k = 1, 2$, we write

$$
\begin{aligned}
\tilde{S}_{\ell,k} &= \frac{1}{q^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\},\pi_J(a_{i,J})=B_k\}\\
&\quad\times\nu^*[\tilde{\pi}_{j_1}(a_{i,j_1});\dots;\tilde{\pi}_{j_{|u|}}(a_{i,j_{|u|}})],\\
S_{\ell,k} &= \frac{1}{q^2\sigma_\ell}\sum_{i=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\},\pi_J(a_{i,J})=B_k\}\\
&\quad\times\nu^*[\pi_{j_1}(a_{i,j_1});\dots;\pi_{j_{|u|}}(a_{i,j_{|u|}})]. \qquad (48)
\end{aligned}
$$

Then for $i = 1, \dots, d-2$,

$$
\begin{aligned}
\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)^2] &= \frac{d(q-1)}{4(i+2)}E[(\tilde{S}_{i,1}+\tilde{S}_{i,2}-S_{i,1}-S_{i,2})^2]\\
&= \frac{d(q-1)}{4(i+2)}E(\tilde{S}_{i,1}^2+\tilde{S}_{i,2}^2+S_{i,1}^2+S_{i,2}^2+2\tilde{S}_{i,1}\tilde{S}_{i,2}+2S_{i,1}S_{i,2}\\
&\quad -2\tilde{S}_{i,1}S_{i,1}-2\tilde{S}_{i,1}S_{i,2}-2\tilde{S}_{i,2}S_{i,1}-2\tilde{S}_{i,2}S_{i,2})\\
&= \frac{d(q-1)}{4(i+2)}E(4S_{i,1}^2+4S_{i,1}S_{i,2}-4\tilde{S}_{i,1}S_{i,1}-4\tilde{S}_{i,1}S_{i,2}). \qquad (49)
\end{aligned}
$$

As in Proposition 1, we have for $i = 1, \dots, d-2$,

$$
\begin{aligned}
\sigma_i^2 &= \frac{1}{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=i+2}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&\quad+\sum_{0\le u_1,\dots,u_d\le 1:|u|=i+2}\frac{(-1)^{|u|-1}|u|}{q^2(q-1)^{|u|-2}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots,\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&\quad+\sum_{0\le u_1,\dots,u_d\le 1:|u|=i+2}\frac{(-1)^{|u|}(q+1-|u|)}{q^2(q-1)^{|u|-1}}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&= \frac{1}{q^2}[1+O(\frac{1}{q})]\sum_{0\le u_1,\cdots,u_d\le 1:|u|=i+2}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},
\end{aligned}
$$

as $q \to \infty$. Hence it follows from (49) and Lemma 5 that

$$
\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)^2] = 1 + O(1/q), \qquad \forall i = 1, \dots, d-2, \qquad (50)
$$

as $q \to \infty$. Finally, (47) is an immediate consequence of (50) and Lemma 1. □

Lemma 5. *Let $S_{\ell,k}$ and $\tilde{S}_{\ell,k}$, $\ell = 1, \dots, d-2$, $k = 1, 2$, be as in (48). Then*

$$
\begin{aligned}
&E(\tilde{S}_{\ell,k}^2) = E(S_{\ell,k}^2)\\
&= \frac{\ell+2}{dq^3\sigma_\ell^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}
\end{aligned}
$$

$$
\begin{aligned}
& +\frac{(-1)^{\ell-1}(\ell+2)}{dq^3(q-1)^{\ell}\sigma_\ell^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},\\
& E(\tilde{S}_{\ell,1}\tilde{S}_{\ell,2}) = E(S_{\ell,1}S_{\ell,2})\\
= \ & \frac{(-1)^{\ell-1}\ell(\ell+2)}{dq^2(q-1)^{\ell+2}\sigma_\ell^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},\\
& E(\tilde{S}_{\ell,1}S_{\ell,2}) = E(S_{\ell,1}\tilde{S}_{\ell,2})\\
= \ & \frac{(-1)^{\ell}\ell(\ell+2)}{dq^2(q-1)^{\ell+1}\sigma_\ell^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},\\
& E(\tilde{S}_{\ell,1}S_{\ell,1}) = E(\tilde{S}_{\ell,2}S_{\ell,2})\\
= \ & -\frac{\ell+2}{dq^3(q-1)\sigma_\ell^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
& +\frac{(-1)^{\ell}(\ell+2)}{dq^3(q-1)^{\ell+1}\sigma_\ell^2} \sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}.
\end{aligned}
$$

PROOF. We observe that for $\ell = 1,\dots,d-2$ and $k = 1,2$,

$$
\begin{aligned}
& E(S_{\ell,k}^2)\\
= \ & \frac{1}{q^4\sigma_\ell^2} E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} \mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\}\}\\
& \times\mathcal{I}\{\pi_J(a_{i_1,J}) = \pi_J(a_{i_2,J}) = B_k\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
& \times\nu^*[\pi_{j_1}(a_{i_2,j_1});\dots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\}\\
= \ & \frac{1}{q^4\sigma_\ell^2} E\Big\{\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} \mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\}\}\\
& \times\mathcal{I}\{\pi_J(a_{i_1,J}) = B_k\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\Big\}\\
& +\frac{1}{q^4\sigma_\ell^2} E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2\ne i_1}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2} \mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\}\}\\
& \times\mathcal{I}\{\pi_J(a_{i_1,J}) = \pi_J(a_{i_2,J}) = B_k\}(-\frac{1}{q-1})^{|u|-1}\\
& \times\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\Big\}\\
= \ & \frac{1}{dq^5\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}\\
& \times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}
\end{aligned}
$$

$$
\begin{aligned}
& +\frac{1}{dq^5\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\mathcal{I}\{a_{i_1,k}=a_{i_2,k}\}(-\frac{1}{q-1})^{|u|-1}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
= \; & \frac{\ell+2}{dq^3\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} \\
& +\frac{\ell+2}{dq^3(q-1)\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}(-\frac{1}{q-1})^{|u|-3} \\
& \quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}, \\
& E(S_{\ell,1}S_{\ell,2}) \\
= \; & \frac{1}{q^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{J\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\mathcal{I}\{\pi_J(a_{i_1,J})=B_1\neq\pi_J(a_{i_2,J})=B_2\} \\
& \quad\times\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
= \; & \frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\mathcal{I}\{a_{i_1,k}\neq a_{i_2,k}\}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \quad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\} \\
= \; & \frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\sum_{1\le l\le|u|:j_l\neq k}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \quad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\} \\
& +\frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\mathcal{I}\{a_{i_1,j_l}\neq a_{i_2,j_l},\forall 1\le l\le|u|\}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \quad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\} \\
= \; & \frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \quad\times\sum_{1\le l\le|u|:j_l\neq k}\mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}(-\frac{1}{q-1})^{|u|-1}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\}\\
&\quad\times\mathcal{I}\{a_{i_1,j_l}\neq a_{i_2,j_l},\forall 1\le l\le |u|\}(-\frac{1}{q-1})^{|u|}E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}\\
=\;&\frac{1}{dq^5\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}|u|(|u|-1)(-\frac{1}{q-1})^{|u|-1}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&+\frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}[q^2-1-|u|(q-1)]\\
&\quad\times|u|(-\frac{1}{q-1})^{|u|}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
=\;&\frac{1}{dq^3(q-1)^2\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}|u|(|u|-1)(-\frac{1}{q-1})^{|u|-3}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&+\frac{1}{dq^3\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}(q+1-|u|)|u|(-\frac{1}{q-1})^{|u|}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
=\;&\frac{1}{dq^3(q-1)^2\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}|u|(|u|-1)(-\frac{1}{q-1})^{|u|-3}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
&-\frac{1}{dq^3(q-1)^2\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}(1-\frac{|u|-2}{q-1})|u|(-\frac{1}{q-1})^{|u|-3}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}\\
=\;&\frac{1}{dq^3(q-1)^2\sigma_\ell^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}(|u|^2-2|u|+\frac{|u|(|u|-2)}{q-1})\\
&\quad\times(-\frac{1}{q-1})^{|u|-3}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},\\
&E(\tilde{S}_{\ell,1}S_{\ell,2})\\
=\;&\frac{1}{q^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{J\in\{j_1,\ldots,j_{|u|}\}\}\\
&\quad\times\mathcal{I}\{\pi_J(a_{i_1,J})=B_1\}\mathcal{I}\{\pi_J(a_{i_2,J})=B_2\}\\
&\quad\times\nu^*[\tilde{\pi}_{j_1}(a_{i_1,j_1});\ldots;\tilde{\pi}_{j_{|u|}}(a_{i_1,j_{|u|}})]\nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\}
\end{aligned}
$$

$$
\begin{aligned}
= \quad & \frac{1}{dq^4\sigma_\ell^2} E\Big\{ \sum_{i_1=1}^{q^2} \sum_{i_2\neq i_1} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} \sum_{k=1}^{d} \mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \times \mathcal{I}\{\pi_k(a_{i_1,k})=B_1\neq \pi_k(a_{i_2,k})=B_2\} \\
& \times \nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \times \nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
= \quad & \frac{1}{dq^5(q-1)\sigma_\ell^2} E\Big\{ \sum_{i_1=1}^{q^2} \sum_{i_2\neq i_1} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} \sum_{k=1}^{d} \mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \times \mathcal{I}\{a_{i_1,k}\neq a_{i_2,k}\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_k(a_{i_2,k});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \times \nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
= \quad & \frac{1}{dq^5(q-1)\sigma_\ell^2} E\Big\{ \sum_{i_1=1}^{q^2} \sum_{i_2\neq i_1} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} \sum_{k=1}^{d} \mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \times \sum_{1\le l\le |u|:j_l\neq k} \mathcal{I}\{a_{i_1,j_l}=a_{i_2,j_l}\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_k(a_{i_2,k});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \times \nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
& + \frac{1}{dq^5(q-1)\sigma_\ell^2} E\Big\{ \sum_{i_1=1}^{q^2} \sum_{i_2\neq i_1} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} \sum_{k=1}^{d} \mathcal{I}\{k\in\{j_1,\ldots,j_{|u|}\}\} \\
& \times \mathcal{I}\{a_{i_1,j_l}\neq a_{i_2,j_l},\forall 1\le l\le |u|\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_k(a_{i_2,k});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})] \\
& \times \nu^*[\pi_{j_1}(a_{i_2,j_1});\ldots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\} \\
= \quad & \frac{1}{dq^5(q-1)\sigma_\ell^2} \sum_{i_1=1}^{q^2} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} |u|(|u|-1)(q-1) \\
& \times \Big(-\frac{1}{q-1}\Big)^{|u|-2} E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
& + \frac{1}{dq^5(q-1)\sigma_\ell^2} \sum_{i_1=1}^{q^2} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} |u|[q^2-1-|u|(q-1)] \\
& \times \Big(-\frac{1}{q-1}\Big)^{|u|-1} E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\ldots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\} \\
= \quad & -\frac{1}{dq^3(q-1)\sigma_\ell^2} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} |u|(|u|-1)\Big(-\frac{1}{q-1}\Big)^{|u|-3} \\
& \times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\} \\
& + \frac{1}{dq^3(q-1)\sigma_\ell^2} \sum_{0\le u_1,\ldots,u_d\le 1:|u|=\ell+2} |u|\Big(1-\frac{|u|-2}{q-1}\Big)\Big(-\frac{1}{q-1}\Big)^{|u|-3} \\
& \times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\ldots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}
\end{aligned}
$$

$$
\begin{aligned}
&= -\frac{1}{dq^3(q-1)\sigma_\ell^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}|u|(|u|-2+\frac{|u|-2}{q-1})\\
&\qquad\times(-\frac{1}{q-1})^{|u|-3}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\},\\
&\quad E(\tilde{S}_{\ell,1}S_{\ell,1})\\
&= \frac{1}{q^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\mathcal{I}\{J\in\{j_1,\dots,j_{|u|}\}\}\\
&\qquad\times\nu^*[\tilde{\pi}_{j_1}(a_{i_1,j_1});\dots;\tilde{\pi}_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\dots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\mathcal{I}\{\pi_J(a_{i_1,J})=\pi_J(a_{i_2,J})=B_1\}\Big\}\\
&= \frac{1}{dq^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\dots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\mathcal{I}\{\pi_k(a_{i_1,k})=\pi_k(a_{i_2,k})=B_1\}\Big\}\\
&= \frac{1}{dq^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}\\
&\qquad\times\mathcal{I}\{\pi_k(a_{i_1,k})=B_1\}\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\Big\}\\
&\quad+\frac{1}{dq^4\sigma_\ell^2}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2\ne i_1}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}\\
&\qquad\times\mathcal{I}\{\pi_k(a_{i_1,k})=\pi_k(a_{i_2,k})=B_1\}\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]\\
&\qquad\times\nu^*[\pi_{j_1}(a_{i_2,j_1});\dots;\pi_{j_{|u|}}(a_{i_2,j_{|u|}})]\Big\}\\
&= \frac{1}{dq^5\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}(-\frac{1}{q-1})\\
&\qquad\times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}\\
&\quad+\frac{1}{dq^4\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2\ne i_1}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_1,\dots,j_{|u|}\}\}\\
&\qquad\times\mathcal{I}\{\pi_k(a_{i_1,k})=\pi_k(a_{i_2,k})=B_1\}(-\frac{1}{q-1})^{|u|}\\
&\qquad\times E\{\nu^*[\pi_{j_1}(a_{i_1,j_1});\dots;\pi_{j_{|u|}}(a_{i_1,j_{|u|}})]^2\}\\
&= -\frac{\ell+2}{dq^3(q-1)\sigma_\ell^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}
\end{aligned}
$$

$$
\begin{aligned}
&-\frac{\ell+2}{dq^3(q-1)^2\sigma_\ell^2}\sum_{0\le u_1,\dots,u_d\le 1:|u|=\ell+2}(-\frac{1}{q-1})^{|u|-3}\\
&\quad\times E\{\nu^*[\pi_{j_1}(a_{1,j_1});\dots;\pi_{j_{|u|}}(a_{1,j_{|u|}})]^2\}.
\end{aligned}
$$

This proves Lemma 5. □

Proposition 5. *Let S_i and $\tilde{S}_i$, $i=1,\dots,d-2$, be as in (37). Then for $i=1,\dots,d-2$,*

$$
\frac{d(q-1)}{4(i+2)}\Big|E[(\tilde{S}_i-S_i)^2\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]-E[(\tilde{S}_i-S_i)^2]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]\Big|\le O(\frac{\|h\|_\infty}{q^{1/2}})\log(1/\varepsilon),
$$

as $q\to\infty$ uniformly over $0<\varepsilon<1$.

Proof. Let $\tilde{S}_{i,k}$ and $S_{i,k}$, $i=1,\dots,d-2$, $k=1,2$, be as in (48). Then it follows from Lemma 5 that

$$
\begin{aligned}
&\frac{d(q-1)}{4(i+2)}\Big|E[(\tilde{S}_i-S_i)^2\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]-E[(\tilde{S}_i-S_i)^2]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]\Big|\\
=\;&\frac{d(q-1)}{4(i+2)}|E\{[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]E^{\mathcal{W}}[(\tilde{S}_i-S_i)^2-E(\tilde{S}_i-S_i)^2]\}|\\
\le\;&\frac{d(q-1)}{4(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(v)|\}E\Big|E^{\mathcal{W}}[(\tilde{S}_{i,1}+\tilde{S}_{i,2}-S_{i,1}-S_{i,2})^2\\
&\quad-E(\tilde{S}_{i,1}+\tilde{S}_{i,2}-S_{i,1}-S_{i,2})^2]\Big|\\
\le\;&\frac{d(q-1)}{4(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(v)|\}\Big\{E|E^{\mathcal{W}}[\tilde{S}_{i,1}^2-E(\tilde{S}_{i,1}^2)]|\\
&\quad+E|E^{\mathcal{W}}[\tilde{S}_{i,2}^2-E(\tilde{S}_{i,2}^2)]|+E|E^{\mathcal{W}}[S_{i,1}^2-E(S_{i,1}^2)]|+E|E^{\mathcal{W}}[S_{i,2}^2-E(S_{i,2}^2)]|\\
&\quad+2E|E^{\mathcal{W}}[\tilde{S}_{i,1}\tilde{S}_{i,2}-E(\tilde{S}_{i,1}\tilde{S}_{i,2})]|+2E|E^{\mathcal{W}}[\tilde{S}_{i,1}S_{i,1}-E(\tilde{S}_{i,1}S_{i,1})]|\\
&\quad+2E|E^{\mathcal{W}}[\tilde{S}_{i,1}S_{i,2}-E(\tilde{S}_{i,1}S_{i,2})]|+2E|E^{\mathcal{W}}[\tilde{S}_{i,2}S_{i,1}-E(\tilde{S}_{i,2}S_{i,1})]|\\
&\quad+2E|E^{\mathcal{W}}[\tilde{S}_{i,2}S_{i,2}-E(\tilde{S}_{i,2}S_{i,2})]|+2E|E^{\mathcal{W}}[S_{i,1}S_{i,2}-E(S_{i,1}S_{i,2})]|\Big\}\\
\le\;&\frac{d(q-1)}{2(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(v)|\}\Big\{E|E^{\mathcal{W}}[\tilde{S}_{i,1}^2-E(\tilde{S}_{i,1}^2)]|\\
&\quad+E|E^{\mathcal{W}}[S_{i,1}^2-E(S_{i,1}^2)]|+E|E^{\mathcal{W}}(\tilde{S}_{i,1}\tilde{S}_{i,2})|+2E|E^{\mathcal{W}}(\tilde{S}_{i,1}S_{i,1})|\\
&\quad+2E|E^{\mathcal{W}}(\tilde{S}_{i,1}S_{i,2})|+E|E^{\mathcal{W}}(S_{i,1}S_{i,2})|+O(q^{-2})\Big\},
\end{aligned}\tag{51}
$$

as $q\to\infty$ uniformly over $0<\varepsilon<1$. Finally we conclude from Lemma 1 and Lemmas 6 to 10 that

$$
\frac{d(q-1)}{4(i+2)}\Big|E[(\tilde{S}_i-S_i)^2\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]-E[(\tilde{S}_i-S_i)^2]E[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]\Big|\le O(\frac{\|h\|_\infty}{q^{1/2}})\log(1/\varepsilon),
$$

as $q\to\infty$ uniformly over $0<\varepsilon<1$. This proves Proposition 5. □

Lemma 6. *With the notation of (51), for* $\ell = 1, \ldots, d-2$,

$$\begin{aligned}
\frac{d(q-1)}{2(\ell+2)} E|E^{\mathcal{W}}[S^2_{\ell,1} - E(S^2_{\ell,1})]| &= O(q^{-1/2}),\\
\frac{d(q-1)}{2(\ell+2)} E|E^{\mathcal{W}}[\tilde{S}^2_{\ell,1} - E(\tilde{S}^2_{\ell,1})]| &= O(q^{-1/2}), \qquad \text{as } q \to \infty.
\end{aligned}$$

Proof. First we observe from (48) that for $\ell = 1, \ldots, d-2$,

$$\begin{aligned}
&E^{\mathcal{W}}(S^2_{\ell,1})\\
&= E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma^2_\ell}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\ \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\\
&\quad\times\mathcal{I}\{J\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J}) = B_1 = \pi_J(a_{i_2,J})\}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}\\
&= \frac{1}{dq^5\sigma^2_\ell}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\ \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\\
&\quad\times\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k} = a_{i_2,k}\}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})].
\end{aligned}$$

Here for $l = 1, \ldots, 4$, given $u^{(l)} = (u_1^{(l)}, \ldots, u_d^{(l)})'$, we write $k \in \{j_{l,1}, \ldots, j_{l,|u^{(l)}|}\}$ if and only if $u_k^{(l)} \ge 1$. Hence

$$\begin{aligned}
&E\{[E^{\mathcal{W}}(S^2_{\ell,1})]^2\}\\
&= E\Big\{\frac{1}{d^2q^{10}\sigma^4_\ell}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\qquad\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\qquad\times\sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\qquad\times\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}\\
&\qquad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\qquad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= R_{\{1,2,3,4\}} + R_{\{1,2,3\},\{4\}} + R_{\{1,2,4\},\{3\}} + R_{\{1,3,4\},\{2\}} + R_{\{2,3,4\},\{1\}}\\
&\quad + R_{\{1,2\},\{3\},\{4\}} + R_{\{1,3\},\{2\},\{4\}} + R_{\{1,4\},\{2\},\{3\}} + R_{\{2,3\},\{1\},\{4\}} + R_{\{2,4\},\{1\},\{3\}}\\
&\quad + R_{\{3,4\},\{1\},\{2\}} + R_{\{1,2\},\{3,4\}} + R_{\{1,3\},\{2,4\}} + R_{\{1,4\},\{2,3\}} + R_{\{1\},\{2\},\{3\},\{4\}}, \qquad (52)
\end{aligned}$$

where given a partition, say $A_1, \ldots, A_p$, of $\{1,2,3,4\}$, (that is $\cup_{i=1}^p A_i = \{1,2,3,4\}$ and $A_i \cap A_j = \emptyset$ whenever $i \neq j$), we define

$$\begin{aligned}
&R_{A_1,\ldots,A_p}\\
=\;& E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum\nolimits^* \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\\
&\times \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\;\sum_{k_1=1}^d\sum_{k_3=1}^d \mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\times\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\},
\end{aligned}\tag{53}$$

and Σ^* denotes summation over $1 \le i_1, i_2, i_3, i_4 \le q^2$ such that if $k, l \in A_j$, then $i_k = i_l$, and if k, l are in different A_j's, then $i_k \neq i_l$. In order to evaluate the terms on the right hand side of (52), it is convenient to further define the following subsets of $\{1, \ldots, d\}$: for $\{l_1, l_2, l_3, l_4\} = \{1,2,3,4\}$,

$$\begin{aligned}
\Theta_{\{1,2,3,4\}} &= \Big\{l \in \cap_{\alpha=1}^4 \{j_{l_\alpha,1},\ldots,j_{l_\alpha,|u^{(l_\alpha)}|}\}\Big\},\\
\Theta_{\{l_1,l_2,l_3\}} &= \Big\{l \in \cap_{\alpha=1}^3 \{j_{l_\alpha,1},\ldots,j_{l_\alpha,|u^{(l_\alpha)}|}\}\Big\}\backslash\{j_{l_4,1},\ldots,j_{l_4,|u^{(l_4)}|}\},\\
\Theta_{\{l_1,l_2\}} &= \Big\{l \in \cap_{\alpha=1}^2 \{j_{l_\alpha,1},\ldots,j_{l_\alpha,|u^{(l_\alpha)}|}\}\Big\}\backslash \cup_{\beta=3}^4 \{j_{l_\beta,1},\ldots,j_{l_\beta,|u^{(l_\beta)}|}\},\\
\Theta_{\{l_1\}} &= \Big\{l \in \{j_{l_1,1},\ldots,j_{l_1,|u^{(l_1)}|}\}\Big\}\backslash \cup_{\beta=2}^4 \{j_{l_\beta,1},\ldots,j_{l_\beta,|u^{(l_\beta)}|}\}.
\end{aligned}$$

Now we observe from Lemma 4 that as $q \to \infty$,

$$\begin{aligned}
&R_{\{1,2,3,4\}}\\
=\;& E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=i_2=i_3=i_4=1}^{q^2}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k_1=1}^d\sum_{k_3=1}^d \mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\times\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}
\end{aligned}$$

$$
\begin{aligned}
&= O(\frac{1}{q^4}),\\
&\quad R_{\{1,2,3\},\{4\}}\\
&= R_{\{1,2,4\},\{3\}} \;=\; R_{\{1,3,4\},\{2\}} \;=\; R_{\{2,3,4\},\{1\}}\\
&= E\Big\{\frac{1}{d^2 q^{10}\sigma_\ell^4} \sum_{i_1=i_2=i_3=1}^{q^2} \sum_{1\le i_4\le q^2: i_4\ne i_1} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1: |u^{(1)}|=\ell+2}\\
&\quad \times \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1: |u^{(2)}|=\ell+2} \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1: |u^{(3)}|=\ell+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1: |u^{(4)}|=\ell+2}\\
&\quad \times \sum_{k_1=1}^{d} \sum_{k_3=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\quad \times \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}\\
&\quad \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= O(\frac{1}{q^3}),\\
&\quad R_{\{1,2\},\{3\},\{4\}} \;=\; R_{\{3,4\},\{1\},\{2\}}\\
&= E\Big\{\frac{1}{d^2 q^{10}\sigma_\ell^4} \sum_{i_1=i_2=1}^{q^2} \sum_{1\le i_3\le q^2: i_3\ne i_1} \sum_{1\le i_4\le q^2: i_4\ne i_1,i_3} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1: |u^{(1)}|=\ell+2}\\
&\quad \times \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1: |u^{(2)}|=\ell+2} \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1: |u^{(3)}|=\ell+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1: |u^{(4)}|=\ell+2}\\
&\quad \times \sum_{k_1=1}^{d} \sum_{k_3=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\quad \times \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}\\
&\quad \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= \frac{O(1)}{q^{10}\sigma_\ell^4} \sum_{i_1=1}^{q^2} \sum_{1\le i_3\le q^2: i_3\ne i_1} \sum_{1\le i_4\le q^2: i_4\ne i_1,i_3} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1: |u^{(1)}|=\ell+2} \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1: |u^{(2)}|=\ell+2}\\
&\quad \times \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1: |u^{(3)}|=\ell+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1: |u^{(4)}|=\ell+2}\\
&\quad \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}0^{|\Theta_{\{4\}}|}
\end{aligned}
$$

$$\begin{aligned}
& \times(\frac{1}{q})^{|\Theta_{\{1,2,3,4\}}|+|\Theta_{\{1,2,4\}}|+|\Theta_{\{1,3,4\}}|+|\Theta_{\{2,3,4\}}|+|\Theta_{\{1,4\}}|+|\Theta_{\{2,4\}}|+|\Theta_{\{3,4\}}|-1} \\
= \; & \frac{O(1)}{q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_3\le q^2:i_3\neq i_1}\sum_{1\le i_4\le q^2:i_4\neq i_1,i_3}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2} \\
& \times\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}(\frac{1}{q})^{|u^{(4)}|-1} \\
= \; & O(\frac{1}{q^3}),
\end{aligned}$$

as $q\to\infty$. In a similar manner, we have

$$\begin{aligned}
& R_{\{1,3\},\{2\},\{4\}} \;=\; R_{\{1,4\},\{2\},\{3\}} \;=\; R_{\{2,3\},\{1\},\{4\}} \;=\; R_{\{2,4\},\{1\},\{3\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=i_3=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\neq i_1}\sum_{1\le i_4\le q^2:i_4\neq i_1,i_2}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times\sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
& \times\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^3}), \\
& R_{\{1,3\},\{2,4\}} \;=\; R_{\{1,4\},\{2,3\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=i_3=1}^{q^2}\sum_{1\le i_2=i_4\le q^2:i_2\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times\sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
& \times\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]
\end{aligned}$$

$$
\begin{aligned}
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^3}).
\end{aligned}
$$

Next we have

$$
\begin{aligned}
&R_{\{1\},\{2\},\{3\},\{4\}} \\
= \;& E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3} \\
&\times\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2} \\
&\times\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
&\times\sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
&\times\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\} \\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& \frac{O(1)}{q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
&\times\sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
&\times\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\} \\
&\times 0^{|\Theta_{\{1\}}|+|\Theta_{\{2\}}|+|\Theta_{\{3\}}|+|\Theta_{\{4\}}|}(\frac{1}{q^2})^{|\Theta_{\{1,2,3,4\}}|+|\Theta_{\{1,2,3\}}|+|\Theta_{\{1,2,4\}}|+|\Theta_{\{1,3,4\}}|+|\Theta_{\{2,3,4\}}|} \\
&\times(\frac{1}{q})^{|\Theta_{\{1,2\}}|+|\Theta_{\{1,3\}}|+|\Theta_{\{1,4\}}|+|\Theta_{\{2,3\}}|+|\Theta_{\{2,4\}}|+|\Theta_{\{3,4\}}|-2}\mathcal{I}\{|\Theta_{\{1,2\}}|\ge|\Theta_{\{3,4\}}|\} \\
= \;& \frac{O(1)}{q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}
\end{aligned}
$$

$$\begin{aligned}
& \times \sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}(\frac{1}{q})^{|u^{(3)}|+|u^{(4)}|-2} \\
= \quad & O(\frac{1}{q^4}).
\end{aligned}$$

The second last equality can be obtained using the heuristic that $a_{i_1,j} \neq a_{i_2,j}$ when $i_1 \neq i_2$ etc. and $k_1 \neq k_3$. However the above bound remains valid when $k_1 = k_3$ or when $a_{i_1,j} = a_{i_2,j}$ since this additional constraint introduces a factor of q while reduces the number of $i_1, \dots, i_4$ by a factor of $1/q$. Finally,

$$\begin{aligned}
& R_{\{1,2\},\{3,4\}} \\
= \quad & E\Big\{\frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=i_2=1}^{q^2}\sum_{1\le i_3=i_4\le q^2:i_4\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times \sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1 \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
& \times\mathcal{I}\{k_3 \in \{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & \frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_3\le q^2:i_3\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times \sum_{k_1=1}^{d}\sum_{k_3=1}^{d}\mathcal{I}\{k_1 \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
& \times\mathcal{I}\{k_3 \in \{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\} \\
& \times E\Big\{\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_3,j_{4,1}});\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_3,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & \frac{1}{d^2q^{10}\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_3\le q^2:i_3\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}|u^{(1)}||u^{(3)}| \\
& \times E\Big\{\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]^2
\end{aligned}$$

$$\begin{aligned}
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]^2\Big\} + O(\frac{1}{q^3}) \\
= \;& \frac{1}{d^2q^6\sigma_\ell^4}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}|u^{(1)}||u^{(3)}| \\
&\times E\{\nu^*[\pi_{j_{1,1}}(a_{1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{1,j_{1,|u^{(1)}|}})]^2\} \\
&\times E\{\nu^*[\pi_{j_{3,1}}(a_{1,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{1,j_{3,|u^{(3)}|}})]^2\} + O(\frac{1}{q^3}) \\
= \;& [E(S_{\ell,1}^2)]^2 + O(\frac{1}{q^3}),
\end{aligned}$$

as $q \to \infty$. The last equality uses Lemma 5. Consequently it follows from (52) that

$$\begin{aligned}
\frac{d(q-1)}{2(\ell+2)}E|E^{\mathcal{W}}[S_{\ell,1}^2 - E(S_{\ell,1}^2)]| &\le \frac{d(q-1)}{2(\ell+2)}\Big\{E\{[E^{\mathcal{W}}(S_{\ell,1}^2) - E(S_{\ell,1}^2)]^2\}\Big\}^{1/2} \\
&= \frac{d(q-1)}{2(\ell+2)}\Big\{E\{[E^{\mathcal{W}}(S_{\ell,1}^2)]^2\} - [E(S_{\ell,1}^2)]^2\Big\}^{1/2} \\
&= O(\frac{1}{q^{1/2}}),
\end{aligned}$$

as $q \to \infty$. Next we observe that

$$\begin{aligned}
& E^{\mathcal{W}}(\tilde{S}_{\ell,1}^2) \\
= \;& E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2} \\
&\times\mathcal{I}\{J \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J}) = B_1 = \pi_J(a_{i_2,J})\} \\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tau_{B_1,B_2}\circ\pi_J(a_{i_1,J});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tau_{B_1,B_2}\circ\pi_J(a_{i_2,J});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \\
= \;& \frac{1}{dq^5(q-1)\sigma_\ell^2}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2} \\
&\times\sum_{k=1}^{d}\mathcal{I}\{k \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k} = a_{i_2,k}\} \\
&\times\sum_{0\le\tilde{c}_k\le q-1:\tilde{c}_k\ne\pi_k(a_{i_1,k})}\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_k;\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_k;\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}$$

and

$$E\{[E^{\mathcal{W}}(\tilde{S}_{\ell,1}^2)]^2\}$$

$$\begin{aligned}
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\quad\times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\quad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\quad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\quad\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\quad\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= \tilde{R}_{\{1,2,3,4\}}+\tilde{R}_{\{1,2,3\},\{4\}}+\tilde{R}_{\{1,2,4\},\{3\}}+\tilde{R}_{\{1,3,4\},\{2\}}+\tilde{R}_{\{2,3,4\},\{1\}}\\
&\quad+\tilde{R}_{\{1,2\},\{3\},\{4\}}+\tilde{R}_{\{1,3\},\{2\},\{4\}}+\tilde{R}_{\{1,4\},\{2\},\{3\}}+\tilde{R}_{\{2,3\},\{1\},\{4\}}+\tilde{R}_{\{2,4\},\{1\},\{3\}}\\
&\quad+\tilde{R}_{\{3,4\},\{1\},\{2\}}+\tilde{R}_{\{1,2\},\{3,4\}}+\tilde{R}_{\{1,3\},\{2,4\}}+\tilde{R}_{\{1,4\},\{2,3\}}+\tilde{R}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}$$

where given a partition, say $A_1,\dots,A_p$, of $\{1,2,3,4\}$, we define

$$\begin{aligned}
\tilde{R}_{A_1,\dots,A_p} &= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum\nolimits^{*}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\\
&\quad\times\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\quad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\quad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\quad\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]
\end{aligned}$$

$$\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\},$$

and Σ^* is as in (53). In a similar way, we observe that

$$\begin{aligned}
&\tilde{R}_{\{1,2,3,4\}}\\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_2=i_3=i_4=1}^{q^2}\ \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \;& O(\frac{1}{q^4}),\\
&\tilde{R}_{\{1,2,3\},\{4\}} \;=\; \tilde{R}_{\{1,2,4\},\{3\}} \;=\; \tilde{R}_{\{1,3,4\},\{2\}} \;=\; \tilde{R}_{\{2,3,4\},\{1\}}\\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_2=i_3=1}^{q^2}\ \sum_{1\le i_4\le q^2:i_4\ne i_1}\ \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]
\end{aligned}$$

$$\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}$$

$$= O(\frac{1}{q^3}),$$

$$\tilde{R}_{\{1,2\},\{3,4\}}$$

$$= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_2=1}^{q^2}\sum_{1\le i_3=i_4\le q^2:i_3\neq i_1}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}$$

$$\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}$$

$$\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}$$

$$\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}$$

$$\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq\pi_{k_1}(a_{i_1,k_1})}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq\pi_{k_3}(a_{i_3,k_3})}$$

$$\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]$$

$$\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]$$

$$\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]$$

$$\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}$$

$$= [E(\tilde{S}_{\ell,1}^2)]^2+O(\frac{1}{q^3}),$$

$$\tilde{R}_{\{1,3\},\{2,4\}} \;=\; \tilde{R}_{\{1,4\},\{2,3\}}$$

$$= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_3=1}^{q^2}\sum_{1\le i_2=i_4\le q^2:i_2\neq i_1}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}$$

$$\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}$$

$$\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}$$

$$\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}$$

$$\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq\pi_{k_1}(a_{i_1,k_1})}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq\pi_{k_3}(a_{i_3,k_3})}$$

$$\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]$$

$$
\begin{aligned}
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \;& O(\frac{1}{q^3}),
\end{aligned}
$$

$$
\begin{aligned}
&\tilde{R}_{\{1,2\},\{3\},\{4\}} \;=\; \tilde{R}_{\{3,4\},\{1\},\{2\}}\\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_2=1}^{q^2}\sum_{1\le i_3\le q^2:i_3\ne i_1}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_3}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \;& O(\frac{1}{q^3}),
\end{aligned}
$$

$$
\begin{aligned}
&\tilde{R}_{\{1\},\{2\},\{3\},\{4\}}\\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}\\
&\times\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}
\end{aligned}
$$

$$
\begin{aligned}
&\times\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq\pi_{k_3}(a_{i_3,k_3})}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
=\ &O(\frac{1}{q^4}),
\end{aligned}
$$

and

$$
\begin{aligned}
&\tilde{R}_{\{1,3\},\{2\},\{4\}}\ =\ \tilde{R}_{\{1,4\},\{2\},\{3\}}\ =\ \tilde{R}_{\{2,3\},\{1\},\{4\}}\ =\ \tilde{R}_{\{2,4\},\{1\},\{3\}}\\
=\ &E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_\ell^4}\sum_{i_1=i_3=1}^{q^2}\ \sum_{1\le i_2\le q^2:i_2\neq i_1}\ \sum_{1\le i_4\le q^2:i_4\neq i_1,i_2}\ \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le1:|u^{(2)}|=\ell+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le1:|u^{(3)}|=\ell+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq\pi_{k_3}(a_{i_3,k_3})}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
=\ &O(\frac{1}{q^4}),
\end{aligned}
$$

as $q\to\infty$. Consequently,

$$
\begin{aligned}
\frac{d(q-1)}{2(\ell+2)}E|E^{\mathcal{W}}[\tilde{S}_{\ell,1}^2-E(\tilde{S}_{\ell,1}^2)]| &\le \frac{d(q-1)}{2(\ell+2)}\Big\{E\{[E^{\mathcal{W}}(\tilde{S}_{\ell,1}^2)-E(\tilde{S}_{\ell,1}^2)]^2\}\Big\}^{1/2}\\
&= \frac{d(q-1)}{2(\ell+2)}\Big\{E\{[E^{\mathcal{W}}(\tilde{S}_{\ell,1}^2)]^2\}-[E(\tilde{S}_{\ell,1}^2)]^2\Big\}^{1/2}\\
&= O(\frac{1}{q^{1/2}}),
\end{aligned}
$$

as $q \to \infty$. This proves Lemma 6. $\square$

Lemma 7. *With the notation of (51), for* $1 \leq \ell_1, \ell_2 \leq d-2$,

$$\frac{d(q-1)}{2(\ell_1+2)} E|E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,2})| = O(\frac{1}{q}), \quad \text{as } q \to \infty.$$

PROOF. We observe that

$$\begin{aligned}
& E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,2}) \\
= \;& E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2} \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2}\; \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \\
& \times \mathcal{I}\{J \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J}) = B_1, \pi_J(a_{i_2,J}) = B_2\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \\
= \;& E^{\mathcal{W}}\Big\{\frac{1}{dq^4\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1} \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2}\; \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \\
& \times \sum_{k=1}^{d}\mathcal{I}\{k \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \mathcal{I}\{\pi_k(a_{i_1,k}) = B_1 \neq \pi_k(a_{i_2,k}) = B_2\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \\
= \;& \frac{1}{dq^5(q-1)\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1} \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2}\; \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2}\; \sum_{k=1}^{d} \\
& \times \mathcal{I}\{k \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k} \neq a_{i_2,k}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}$$

and

$$\begin{aligned}
& E\{[E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,2})]^2\} \\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2} \sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{i_3=1}^{q^2}\sum_{i_4\neq i_3} \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2} \\
& \quad \times \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2}\; \sum_{0\leq u_1^{(3)},\ldots,u_d^{(3)}\leq 1:|u^{(3)}|=\ell_1+2}\; \sum_{0\leq u_1^{(4)},\ldots,u_d^{(4)}\leq 1:|u^{(4)}|=\ell_2+2} \\
& \quad \times \sum_{k_1=1}^{d}\mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}
\end{aligned}$$

$$
\begin{aligned}
&\quad \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\} \\
&\quad \times \mathcal{I}\{a_{i_1,k_1} \neq a_{i_2,k_1}\} \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
&\quad \times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\quad \times \mathcal{I}\{a_{i_3,k_3} \neq a_{i_4,k_3}\} \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
&\quad \times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= &\quad R^{(1,2)}_{\{1,3\},\{2\},\{4\}} + R^{(1,2)}_{\{1,4\},\{2\},\{3\}} + R^{(1,2)}_{\{2,3\},\{1\},\{4\}} + R^{(1,2)}_{\{2,4\},\{1\},\{3\}} \\
&\quad + R^{(1,2)}_{\{1,3\},\{2,4\}} + R^{(1,2)}_{\{1,4\},\{2,3\}} + R^{(1,2)}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
& R^{(1,2)}_{\{1,3\},\{2\},\{4\}} \;=\; R^{(1,2)}_{\{1,4\},\{2\},\{3\}} \;=\; R^{(1,2)}_{\{2,3\},\{1\},\{4\}} \;=\; R^{(1,2)}_{\{2,4\},\{1\},\{3\}} \\
= \;& E\Big\{\frac{1}{d^2 q^{10}(q-1)^2 \sigma^2_{\ell_1} \sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1 \le i_2 \le q^2 : i_2 \neq i_1} \sum_{1 \le i_4 \le q^2 : i_4 \neq i_1, i_2} \sum_{0 \le u_1^{(1)}, \ldots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
& \times \sum_{0 \le u_1^{(2)}, \ldots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \ldots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \ldots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \ldots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \ldots, j_{2,|u^{(2)}|}\}\} \mathcal{I}\{a_{i_1,k_1} \neq a_{i_2,k_1}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\} \mathcal{I}\{a_{i_1,k_3} \neq a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})] \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^5}), \\
& R^{(1,2)}_{\{1,3\},\{2,4\}} \;=\; R^{(1,2)}_{\{1,4\},\{2,3\}} \\
= \;& E\Big\{\frac{1}{d^2 q^{10}(q-1)^2 \sigma^2_{\ell_1} \sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1 \le i_2 \le q^2 : i_2 \neq i_1} \sum_{0 \le u_1^{(1)}, \ldots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
& \times \sum_{0 \le u_1^{(2)}, \ldots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \ldots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \ldots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \ldots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \ldots, j_{2,|u^{(2)}|}\}\} \mathcal{I}\{a_{i_1,k_1} \neq a_{i_2,k_1}\}
\end{aligned}
$$

$$
\begin{aligned}
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3} \neq a_{i_2,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_2,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_2,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^4}), \\
& R^{(1,2)}_{\{1\},\{2\},\{3\},\{4\}} \\
= \; & E\Big\{\frac{1}{d^2 q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1 \le i_2 \le q^2 : i_2 \neq i_1} \sum_{1 \le i_3 \le q^2 : i_3 \neq i_1, i_2} \sum_{1 \le i_4 \le q^2 : i_4 \neq i_1, i_2, i_3} \\
& \times \sum_{0 \le u_1^{(1)}, \ldots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \; \sum_{0 \le u_1^{(2)}, \ldots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \\
& \times \sum_{0 \le u_1^{(3)}, \ldots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \; \sum_{0 \le u_1^{(4)}, \ldots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \ldots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \ldots, j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1} \neq a_{i_2,k_1}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} \neq a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^6}),
\end{aligned}
$$

as $q \to \infty$. Thus we conclude that

$$
\frac{d(q-1)}{2(\ell_1+2)} E|E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,2})| \le \frac{d(q-1)}{2(\ell_1+2)} \{E[E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,2})]^2\}^{1/2} = O(\frac{1}{q}),
$$

as $q \to \infty$. This proves Lemma 7. □

Lemma 8. *With the notation of (51), for* $1 \le \ell_1, \ell_2 \le d-2$,

$$
\frac{d(q-1)}{\ell_1+2} E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,2})| = O(\frac{1}{q}), \qquad \textit{as } q \to \infty.
$$

Proof. We observe that

$$
\begin{aligned}
& E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,2}) \\
= \; & E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{i_2 \neq i_1} \sum_{0 \le u_1^{(1)}, \ldots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \; \sum_{0 \le u_1^{(2)}, \ldots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2}
\end{aligned}
$$

$$
\begin{aligned}
&\times\mathcal{I}\{J\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J})=B_1,\pi_J(a_{i_2,J})=B_2\}\\
&\times\nu^*[\tilde\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}\\
= \quad & E^{\mathcal{W}}\Big\{\frac{1}{dq^4\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{k=1}^{d}\\
&\times\mathcal{I}\{k\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_k(a_{i_1,k})=B_1,\pi_k(a_{i_2,k})=B_2\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}\\
= \quad & \frac{1}{dq^5(q-1)\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
&\quad\times\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k}\neq a_{i_2,k}\}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_k(a_{i_2,k});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}
$$

and

$$
\begin{aligned}
& E\{[E^{\mathcal{W}}(\tilde S_{\ell_1,1}S_{\ell_2,2})]^2\}\\
= \quad & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{i_3=1}^{q^2}\sum_{i_4\neq i_3}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\quad\times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\quad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}\neq a_{i_2,k_1}\}\\
&\quad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}\neq a_{i_4,k_3}\}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{k_1}(a_{i_2,k_1});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{k_3}(a_{i_4,k_3});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\quad\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \quad & R^{(\tilde 1,2)}_{\{1,3\},\{2\},\{4\}}+R^{(\tilde 1,2)}_{\{1,4\},\{2\},\{3\}}+R^{(\tilde 1,2)}_{\{2,3\},\{1\},\{4\}}+R^{(\tilde 1,2)}_{\{2,4\},\{1\},\{3\}}\\
& +R^{(\tilde 1,2)}_{\{1,3\},\{2,4\}}+R^{(\tilde 1,2)}_{\{1,4\},\{2,3\}}+R^{(\tilde 1,2)}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
& R^{(\tilde{1},2)}_{\{1,3\},\{2\},\{4\}} \;=\; R^{(\tilde{1},2)}_{\{1,4\},\{2\},\{3\}} \;=\; R^{(\tilde{1},2)}_{\{2,3\},\{1\},\{4\}} \;=\; R^{(\tilde{1},2)}_{\{2,4\},\{1\},\{3\}} \\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1\le i_2\le q^2:i_2\ne i_1} \sum_{1\le i_4\le q^2:i_4\ne i_2,i_3} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\mathcal{I}\{a_{i_1,k_1}\ne a_{i_2,k_1}\} \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{k_3}(a_{i_4,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})] \\
& \times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\mathcal{I}\{a_{i_1,k_3}\ne a_{i_4,k_3}\}\Big\} \\
= \;& O(\frac{1}{q^4}), \\
& R^{(\tilde{1},2)}_{\{1,3\},\{2,4\}} \;=\; R^{(\tilde{1},2)}_{\{1,4\},\{2,3\}} \\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1\le i_2\le q^2:i_2\ne i_1} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\mathcal{I}\{a_{i_1,k_1}\ne a_{i_2,k_1}\} \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{k_3}(a_{i_2,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})] \\
& \times \nu^*[\pi_{j_{4,1}}(a_{i_2,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_2,j_{4,|u^{(4)}|}})]\mathcal{I}\{a_{i_1,k_3}\ne a_{i_2,k_3}\}\Big\} \\
= \;& O(\frac{1}{q^4}),
\end{aligned}
$$

and

$$\begin{aligned} R^{(\tilde{1},2)}_{\{1\},\{2\},\{3\},\{4\}} &= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\neq i_1}\sum_{1\le i_3\le q^2:i_3\neq i_1,i_2} \\ &\times\sum_{1\le i_4\le q^2:i_4\neq i_1,i_2,i_3}\sum_{0\le u^{(1)}_1,\ldots,u^{(1)}_d\le 1:|u^{(1)}|=\ell_1+2}\sum_{0\le u^{(2)}_1,\ldots,u^{(2)}_d\le 1:|u^{(2)}|=\ell_2+2} \\ &\times\sum_{0\le u^{(3)}_1,\ldots,u^{(3)}_d\le 1:|u^{(3)}|=\ell_1+2}\sum_{0\le u^{(4)}_1,\ldots,u^{(4)}_d\le 1:|u^{(4)}|=\ell_2+2} \\ &\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\ &\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\} \\ &\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\ &\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\mathcal{I}\{a_{i_1,k_1}\neq a_{i_2,k_1}\} \\ &\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{k_3}(a_{i_4,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\ &\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\mathcal{I}\{a_{i_3,k_3}\neq a_{i_4,k_3}\}\Big\} \\ &= O(\frac{1}{q^4}), \end{aligned}$$

as $q\to\infty$. Thus we conclude that

$$\frac{d(q-1)}{\ell_1+2}E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,2})|\le\frac{d(q-1)}{\ell_1+2}\{E[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,2})]^2\}^{1/2}=O(\frac{1}{q}),$$

as $q\to\infty$. This proves Lemma 8. □

Lemma 9. *With the notation of (51), for* $1\le\ell_1,\ell_2\le d-2$,

$$\frac{d(q-1)}{\ell_1+2}E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,1})|=O(\frac{1}{q}),\qquad \textit{as } q\to\infty.$$

Proof. We observe that

$$\begin{aligned} &E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,1}) \\ &= E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u^{(1)}_1,\ldots,u^{(1)}_d\le 1:|u^{(1)}|=\ell_1+2}\sum_{0\le u^{(2)}_1,\ldots,u^{(2)}_d\le 1:|u^{(2)}|=\ell_2+2} \\ &\times\mathcal{I}\{J\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J})=B_1=\pi_J(a_{i_2,J})\} \\ &\times\nu^*[\tilde{\pi}_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{\pi}_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \end{aligned}$$

$$
\begin{aligned}
= {} & E^{\mathcal{W}}\Big\{\frac{1}{dq^4\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{k=1}^{d} \\
& \times\mathcal{I}\{k\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_k(a_{i_1,k})=B_1=\pi_k(a_{i_2,k})\} \\
& \quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tau_{B_1,B_2}\circ\pi_k(a_{i_1,k});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \\
= {} & -\frac{1}{dq^5(q-1)\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \\
& \times\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_2,k}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}
$$

and

$$
\begin{aligned}
& E\{[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,1})]^2\} \\
= {} & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \\
& \times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\} \\
& \times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= {} & R^{(\tilde{1},1)}_{\{1,2,3,4\}}+R^{(\tilde{1},1)}_{\{1,2,3\},\{4\}}+R^{(\tilde{1},1)}_{\{1,2,4\},\{3\}}+R^{(\tilde{1},1)}_{\{1,3,4\},\{2\}}+R^{(\tilde{1},1)}_{\{2,3,4\},\{1\}} \\
& +R^{(\tilde{1},1)}_{\{1,2\},\{3\},\{4\}}+R^{(\tilde{1},1)}_{\{1,3\},\{2\},\{4\}}+R^{(\tilde{1},1)}_{\{1,4\},\{2\},\{3\}}+R^{(\tilde{1},1)}_{\{2,3\},\{1\},\{4\}}+R^{(\tilde{1},1)}_{\{2,4\},\{1\},\{3\}} \\
& +R^{(\tilde{1},1)}_{\{3,4\},\{1\},\{2\}}+R^{(\tilde{1},1)}_{\{1,2\},\{3,4\}}+R^{(\tilde{1},1)}_{\{1,3\},\{2,4\}}+R^{(\tilde{1},1)}_{\{1,4\},\{2,3\}}+R^{(\tilde{1},1)}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
& R^{(\tilde{1},1)}_{\{1,2,3,4\}} \\
= {} & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}
\end{aligned}
$$

$$
\begin{aligned}
& \times \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \ \sum_{0\leq u_1^{(3)},\ldots,u_d^{(3)}\leq 1:|u^{(3)}|=\ell_1+2} \ \sum_{0\leq u_1^{(4)},\ldots,u_d^{(4)}\leq 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_1,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_1,j_{4,|u^{(4)}|}})]\Big\} \\
= \ & O(\frac{1}{q^6}),
\end{aligned}
$$

$$
\begin{aligned}
& R^{(\tilde{1},1)}_{\{1,2,3\},\{4\}} \ = \ R^{(\tilde{1},1)}_{\{1,2,4\},\{3\}} \ = \ R^{(\tilde{1},1)}_{\{1,3,4\},\{2\}} \ = \ R^{(\tilde{1},1)}_{\{2,3,4\},\{1\}} \\
= \ & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2} \sum_{i_1=1}^{q^2} \ \sum_{1\leq i_4\leq q^2:i_4\neq i_1} \ \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \ \sum_{0\leq u_1^{(3)},\ldots,u_d^{(3)}\leq 1:|u^{(3)}|=\ell_1+2} \ \sum_{0\leq u_1^{(4)},\ldots,u_d^{(4)}\leq 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3} = a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \ & O(\frac{1}{q^5}),
\end{aligned}
$$

$$
\begin{aligned}
& R^{(\tilde{1},1)}_{\{1,2\},\{3\},\{4\}} \ = \ R^{(\tilde{1},1)}_{\{3,4\},\{1\},\{2\}} \\
= \ & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}\sigma_{\ell_2}^2} \sum_{i_1=1}^{q^2} \ \sum_{1\leq i_3\leq q^2:i_3\neq i_1} \ \sum_{1\leq i_4\leq q^2:i_4\neq i_1,i_3} \ \sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \ \sum_{0\leq u_1^{(3)},\ldots,u_d^{(3)}\leq 1:|u^{(3)}|=\ell_1+2} \ \sum_{0\leq u_1^{(4)},\ldots,u_d^{(4)}\leq 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}
\end{aligned}
$$

$$
\begin{aligned}
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^5}), \\
& R^{(\tilde{1},1)}_{\{1,3\},\{2\},\{4\}} \;=\; R^{(\tilde{1},1)}_{\{1,4\},\{2\},\{3\}} \;=\; R^{(\tilde{1},1)}_{\{2,3\},\{1\},\{4\}} \;=\; R^{(\tilde{1},1)}_{\{2,4\},\{1\},\{3\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1\le i_2\le q^2 : i_2\ne i_1} \sum_{1\le i_4\le q^2 : i_4\ne i_1, i_2} \sum_{0\le u^{(1)}_1,\ldots,u^{(1)}_d\le 1 : |u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\le u^{(2)}_1,\ldots,u^{(2)}_d\le 1 : |u^{(2)}|=\ell_2+2} \sum_{0\le u^{(3)}_1,\ldots,u^{(3)}_d\le 1 : |u^{(3)}|=\ell_1+2} \sum_{0\le u^{(4)}_1,\ldots,u^{(4)}_d\le 1 : |u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \ldots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \ldots, j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3} = a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^6}), \\
& R^{(\tilde{1},1)}_{\{1,2\},\{3,4\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1\le i_3\le q^2 : i_3\ne i_1} \sum_{0\le u^{(1)}_1,\ldots,u^{(1)}_d\le 1 : |u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\le u^{(2)}_1,\ldots,u^{(2)}_d\le 1 : |u^{(2)}|=\ell_2+2} \sum_{0\le u^{(3)}_1,\ldots,u^{(3)}_d\le 1 : |u^{(3)}|=\ell_1+2} \sum_{0\le u^{(4)}_1,\ldots,u^{(4)}_d\le 1 : |u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \ldots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \ldots, j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \ldots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \ldots, j_{4,|u^{(4)}|}\}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}}); \ldots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \ldots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_3,j_{4,1}}); \ldots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_3,j_{4,|u^{(4)}|}})]\Big\}
\end{aligned}
$$

$$
\begin{aligned}
&= O(\frac{1}{q^4}),\\
&R^{(\tilde{1},1)}_{\{1,3\},\{2,4\}} \;=\; R^{(\tilde{1},1)}_{\{1,4\},\{3,4\}}\\
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3}=a_{i_2,k_3}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_2,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_2,j_{4,|u^{(4)}|}})]\Big\}\\
&= O(\frac{1}{q^5}),\\
&R^{(\tilde{1},1)}_{\{1\},\{2\},\{3\},\{4\}}\\
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}\\
&\times\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
&\times\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= O(\frac{1}{q^6}),
\end{aligned}
$$

as $q\to\infty$. Thus we conclude that

$$
\frac{d(q-1)}{\ell_1+2}E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,1})|\le\frac{d(q-1)}{\ell_1+2}\{E[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}S_{\ell_2,1})]^2\}^{1/2}=O(\frac{1}{q}),
$$

as $q \to \infty$. This proves Lemma 9. $\square$

Lemma 10. *With the notation of (51), for* $1 \le \ell_1, \ell_2 \le d-2$,

$$\frac{d(q-1)}{\ell_1+2} E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,2})| = O(\frac{1}{q}), \qquad \text{as } q \to \infty.$$

Proof. We observe that

$$\begin{aligned}
&E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,2})\\
&= E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
&\quad\times\mathcal{I}\{J\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J})=B_1,\pi_J(a_{i_2,J})=B_2\}\\
&\quad\times\nu^*[\tilde{\pi}_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{\pi}_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\tilde{\pi}_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{\pi}_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}\\
&= \frac{1}{dq^5(q-1)\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
&\qquad\times\sum_{k=1}^{d}\mathcal{I}\{k\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k}\neq a_{i_2,k}\}\\
&\qquad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_k(a_{i_2,k});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\qquad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_k(a_{i_1,k});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}$$

and

$$\begin{aligned}
&E\{[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,2})]^2\}\\
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=1}^{q^2}\sum_{i_2\neq i_1}\sum_{i_3=1}^{q^2}\sum_{i_4\neq i_3}\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\qquad\times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\qquad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}\neq a_{i_2,k_1}\}\\
&\qquad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}\neq a_{i_4,k_3}\}\\
&\qquad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{k_1}(a_{i_2,k_1});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\qquad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{k_1}(a_{i_1,k_1});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\qquad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{k_3}(a_{i_4,k_3});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]
\end{aligned}$$

$$
\begin{aligned}
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{k_3}(a_{i_3,k_3});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \ & \tilde{R}^{(1,2)}_{\{1,3\},\{2\},\{4\}}+\tilde{R}^{(1,2)}_{\{1,4\},\{2\},\{3\}}+\tilde{R}^{(1,2)}_{\{2,3\},\{1\},\{4\}}+\tilde{R}^{(1,2)}_{\{2,4\},\{1\},\{3\}}\\
&+\tilde{R}^{(1,2)}_{\{1,3\},\{2,4\}}+\tilde{R}^{(1,2)}_{\{1,4\},\{2,3\}}+\tilde{R}^{(1,2)}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
&\tilde{R}^{(1,2)}_{\{1,3\},\{2\},\{4\}} \ = \ \tilde{R}^{(1,2)}_{\{1,4\},\{2\},\{3\}} \ = \ \tilde{R}^{(1,2)}_{\{2,3\},\{1\},\{4\}} \ = \ \tilde{R}^{(1,2)}_{\{2,4\},\{1\},\{3\}}\\
= \ & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2}\\
&\times\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
&\times\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}\ne a_{i_2,k_1}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3}\ne a_{i_4,k_3}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{k_1}(a_{i_1,k_1});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{k_3}(a_{i_4,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{k_3}(a_{i_1,k_3});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \ & O(\frac{1}{q^5}),\\
&\tilde{R}^{(1,2)}_{\{1,3\},\{2,4\}} \ = \ \tilde{R}^{(1,2)}_{\{1,4\},\{2,3\}}\\
= \ & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}\ne a_{i_2,k_1}\}\\
&\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3}\ne a_{i_2,k_3}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]
\end{aligned}
$$

$$
\begin{aligned}
& \times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{k_1}(a_{i_1,k_1});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_1,j_{3,1}});\ldots;\pi_{k_3}(a_{i_2,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_1,j_{3,|u^{(3)}|}})] \\
& \times\nu^*[\pi_{j_{4,1}}(a_{i_2,j_{4,1}});\ldots;\pi_{k_3}(a_{i_1,k_3});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_2,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^4}), \\
& \tilde{R}^{(1,2)}_{\{1\},\{2\},\{3\},\{4\}} \\
= \;& E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\;\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2} \\
& \times\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \\
& \times\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}\ne a_{i_2,k_1}\} \\
& \times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}\ne a_{i_4,k_3}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{k_1}(a_{i_2,k_1});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{k_1}(a_{i_1,k_1});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{k_3}(a_{i_4,k_3});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
& \times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{k_3}(a_{i_3,k_3});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^6}),
\end{aligned}
$$

as $q\to\infty$. Thus we conclude that

$$
\frac{d(q-1)}{\ell_1+2}E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,2})|\le\frac{d(q-1)}{\ell_1+2}\{E[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,2})]^2\}^{1/2}=O(\frac{1}{q}),
$$

as $q\to\infty$. This proves Lemma 10. □

Proposition 6. *Let S_i and $\tilde{S}_i$, $i=1,\ldots,d-2$, be as in (37). Then for $1\le i\ne j\le d-2$,*

$$
|\frac{d(q-1)}{4(i+2)}E\{(\tilde{S}_i-S_i)^2\int_0^1[\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V+t(\tilde{V}-V))-\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]dt\}|\le O(\frac{\|h\|_\infty}{\varepsilon q^{1/2}}),
$$

and

$$
\begin{aligned}
& |\frac{d(q-1)}{4(i+2)}E\{(\tilde{S}_i-S_i)(\tilde{S}_j-S_j)\int_0^1[\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V+t(\tilde{V}-V))-\frac{\partial^2}{\partial v_i\partial v_j}\psi_{\varepsilon^2}(V)]dt\}| \\
\le \;& O(\frac{\|h\|_\infty}{\varepsilon q^{1/2}}),
\end{aligned}
$$

as $q \to \infty$ *uniformly over* $0 < \varepsilon < 1$.

Proof. Using Taylor series and Lemma 1, we observe that for $0 < \varepsilon < 1$ and $i = 1, \ldots, d-2$,

$$\begin{aligned}
&|\frac{d(q-1)}{4(i+2)} E\{(\tilde{S}_i - S_i)^2 \int_0^1 [\frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V + t(\tilde{V} - V)) - \frac{\partial^2}{\partial v_i^2}\psi_{\varepsilon^2}(V)]dt\}| \\
\leq\ & \sum_{j=1}^{d-2} \frac{d(q-1)}{4(i+2)} E[(\tilde{S}_i - S_i)^2 |\tilde{S}_j - S_j| \sup_{v \in \mathbb{R}^{d-2}} |\frac{\partial^3}{\partial v_i^2 \partial v_j}\psi_{\varepsilon^2}(v)|] \\
\leq\ & \frac{c\|h\|_\infty}{\varepsilon} \sum_{j=1}^{d-2} \frac{d(q-1)}{4(i+2)} \{E[(\tilde{S}_i - S_i)^4]\}^{1/2} \{E[(\tilde{S}_j - S_j)^2]\}^{1/2} \\
\leq\ & \frac{2^{1/2} c\|h\|_\infty}{\varepsilon} \sum_{j=1}^{d-2} \frac{d(q-1)}{4(i+2)} [E(S_i^4)]^{1/2} \{E[(\tilde{S}_j - S_j)^2]\}^{1/2},
\end{aligned}$$

where c is a constant depending only on d. In a similar fashion, we have for $1 \leq i \neq j \leq d-2$,

$$\begin{aligned}
&|\frac{d(q-1)}{4(i+2)} E\{(\tilde{S}_i - S_i)(\tilde{S}_j - S_j) \int_0^1 [\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V + t(\tilde{V} - V)) - \frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V)]dt\}| \\
\leq\ & \sum_{k=1}^{d-2} \frac{d(q-1)}{4(i+2)} E[|\tilde{S}_i - S_i||\tilde{S}_j - S_j||\tilde{S}_k - S_k| \sup_{v \in \mathbb{R}^{d-2}} |\frac{\partial^3}{\partial v_i \partial v_j \partial v_k}\psi_{\varepsilon^2}(v)|] \\
\leq\ & \frac{c\|h\|_\infty}{\varepsilon} \sum_{k=1}^{d-2} \frac{d(q-1)}{4(i+2)} \{E[(\tilde{S}_i - S_i)^4]\}^{1/4} \{E[(\tilde{S}_j - S_j)^4]\}^{1/4} \{E[(\tilde{S}_k - S_k)^2]\}^{1/2} \\
\leq\ & \frac{2^{1/2} c\|h\|_\infty}{\varepsilon} \sum_{k=1}^{d-2} \frac{d(q-1)}{4(i+2)} [E(S_i^4)]^{1/4} [E(S_j^4)]^{1/4} \{E[(\tilde{S}_k - S_k)^2]\}^{1/2}.
\end{aligned}$$

Proposition 6 now follows from Proposition 4 and Lemma 11. □

Lemma 11. *Let* $S_\ell, \ell = 1, \ldots, d-2$, *be as in (37). Then* $E(S_\ell^4) = O(1/q^2)$ *as* $q \to \infty$.

Proof. We observe from (48) that $S_\ell^4 = S_{\ell,1}^4 + S_{\ell,2}^4$ and hence

$$\begin{aligned}
E(S_\ell^4) &= 2E(S_{\ell,1}^4) \\
= & \frac{2}{q^8 \sigma_\ell^4} E\Big\{ \sum_{i_1=1}^{q^2} \sum_{i_2=1}^{q^2} \sum_{i_3=1}^{q^2} \sum_{i_4=1}^{q^2} \sum_{0 \leq u_1^{(1)}, \ldots, u_d^{(1)} \leq 1 : |u^{(1)}| = \ell+2} \\
& \times \sum_{0 \leq u_1^{(2)}, \ldots, u_d^{(2)} \leq 1 : |u^{(2)}| = \ell+2} \sum_{0 \leq u_1^{(3)}, \ldots, u_d^{(3)} \leq 1 : |u^{(3)}| = \ell+2} \sum_{0 \leq u_1^{(4)}, \ldots, u_d^{(4)} \leq 1 : |u^{(4)}| = \ell+2} \\
& \times \mathcal{I}\{J \in \cap_{l=1}^4 \{j_{l,1}, \ldots, j_{l,|u^{(l)}|}\}\} \mathcal{I}\{B_1 = \pi_J(a_{i_l,J}), l = 1, \ldots, 4\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \ldots; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \ldots; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]
\end{aligned}$$

$$
\begin{aligned}
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
=\;&\frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k=1}^{d}\mathcal{I}\{k\in\cap_{l=1}^4\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_2,k}=a_{i_3,k}=a_{i_4,k}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
=\;&R^*_{\{1,2,3,4\}}+R^*_{\{1,2,3\},\{4\}}+R^*_{\{1,2,4\},\{3\}}+R^*_{\{1,3,4\},\{2\}}\\
&+R^*_{\{2,3,4\},\{1\}}+R^*_{\{1,2\},\{3\},\{4\}}+R^*_{\{1,3\},\{2\},\{4\}}+R^*_{\{1,4\},\{2\},\{3\}}\\
&+R^*_{\{2,3\},\{1\},\{4\}}+R^*_{\{2,4\},\{1\},\{3\}}+R^*_{\{3,4\},\{1\},\{2\}}+R^*_{\{1,2\},\{3,4\}}\\
&+R^*_{\{1,3\},\{2,4\}}+R^*_{\{1,4\},\{2,3\}}+R^*_{\{1\},\{2\},\{3\},\{4\}},\qquad(54)
\end{aligned}
$$

where

$$
\begin{aligned}
&R^*_{\{1,2,3,4\}}\\
=\;&\frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=i_2=i_3=i_4=1}^{q^2}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\;\sum_{k=1}^{d}\mathcal{I}\{k\in\cap_{l=1}^4\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\\
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
=\;&O(\frac{1}{q^3}),\\
&R^*_{\{1,2,3\},\{4\}}\;=\;R^*_{\{1,2,4\},\{3\}}\;=\;R^*_{\{1,3,4\},\{2\}}\;=\;R^*_{\{2,3,4\},\{1\}}\\
=\;&\frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=i_2=i_3=1}^{q^2}\;\sum_{1\le i_4\le q^2:i_4\ne i_1}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\\
&\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2}\\
&\times\sum_{k=1}^{d}\mathcal{I}\{k\in\cap_{l=1}^4\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_4,k}\}
\end{aligned}
$$

$$
\begin{aligned}
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^4}),
\end{aligned}
$$

$$
\begin{aligned}
& R^*_{\{1,2\},\{3,4\}} \;=\; R^*_{\{1,3\},\{2,4\}} \;=\; R^*_{\{1,4\},\{2,3\}} \\
= \;& \frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=i_2=1}^{q^2}\;\sum_{1\le i_3=i_4\le q^2:i_3\ne i_1}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times\sum_{k=1}^{d}\mathcal{I}\{k\in\cap_{l=1}^4\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_3,k}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_3,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_3,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^2}),
\end{aligned}
$$

$$
\begin{aligned}
& R^*_{\{1,2\},\{3\},\{4\}} \\
= \;& \frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=i_2=1}^{q^2}\;\sum_{1\le i_3\le q^2:i_3\ne i_1}\;\sum_{1\le i_4\le q^2:i_4\ne i_1,i_3}\;\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2} \\
& \times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}\;\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2}\;\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times\sum_{k=1}^{d}\mathcal{I}\{k\in\cap_{l=1}^4\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_3,k}=a_{i_4,k}\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})] \\
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^3}),
\end{aligned}
$$

$$
\begin{aligned}
& R^*_{\{1\},\{2\},\{3\},\{4\}} \\
= \;& \frac{2}{dq^9\sigma_\ell^4}E\Big\{\sum_{i_1=1}^{q^2}\;\sum_{1\le i_2\le q^2:i_2\ne i_1}\;\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\;\sum_{1\le i_4\le q^2:i_4\ne i_{1,2},i_3} \\
& \times\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell+2}\;\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell+2}
\end{aligned}
$$

$$
\begin{aligned}
& \times \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell+2} \; \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell+2} \\
& \times \sum_{k=1}^{d} \mathcal{I}\{k \in \cap_{l=1}^{4}\{j_{l,1},\ldots,j_{l,|u^{(l)}|}\}\}\mathcal{I}\{a_{i_1,k}=a_{i_2,k}=a_{i_3,k}=a_{i_4,k}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \;& O(\frac{1}{q^4}),
\end{aligned}
$$

as $q \to \infty$. Consequently we conclude from (54) that $E(S_\ell^4) = O(1/q^2)$ as $q \to \infty$. This proves Lemma 11. □

Proposition 7. *Let S_i and $\tilde{S}_i$, $i = 1,\ldots,d-2$, be as in (37). Then for $1 \le i \ne j \le d-2$,*

$$
|\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)(\tilde{S}_j - S_j)\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V)]| = O(\frac{\|h\|_\infty}{q^{1/2}})\log(1/\varepsilon),
$$

as $q \to \infty$ uniformly over $0 < \varepsilon < 1$.

Proof. Let $S_{i,k}$ and $\tilde{S}_{i,k}$, $i = 1,\ldots,d-2$ and $k = 1,2$, be as in (48). Then $S_i = S_{i,1} + S_{i,2}$ and $\tilde{S}_i = \tilde{S}_{i,1} + \tilde{S}_{i,2}$. For $1 \le i \ne j \le d-2$, we observe that

$$
\begin{aligned}
& |\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)(\tilde{S}_j - S_j)\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V)]| \\
= \;& |\frac{d(q-1)}{4(i+2)}E\{\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V)E^{\mathcal{W}}[(\tilde{S}_i - S_i)(\tilde{S}_j - S_j)]\}| \\
= \;& \frac{d(q-1)}{4(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(v)|\} \\
& \times E|E^{\mathcal{W}}[(\tilde{S}_{i,1} + \tilde{S}_{i,2} - S_{i,1} - S_{i,2})(\tilde{S}_{j,1} + \tilde{S}_{j,2} - S_{j,1} - S_{j,2})]| \\
= \;& \frac{d(q-1)}{4(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(v)|\}E|E^{\mathcal{W}}(\tilde{S}_{i,1}\tilde{S}_{j,1} + \tilde{S}_{i,1}\tilde{S}_{j,2} - \tilde{S}_{i,1}S_{j,1} - \tilde{S}_{i,1}S_{j,2} \\
& + \tilde{S}_{i,2}\tilde{S}_{j,1} + \tilde{S}_{i,2}\tilde{S}_{j,2} - \tilde{S}_{i,2}S_{j,1} - \tilde{S}_{i,2}S_{j,2} - S_{i,1}\tilde{S}_{j,1} - S_{i,1}\tilde{S}_{j,2} + S_{i,1}S_{j,1} + S_{i,1}S_{j,2} \\
& - S_{i,2}\tilde{S}_{j,1} - S_{i,2}\tilde{S}_{j,2} + S_{i,2}S_{j,1} + S_{i,2}S_{j,2})| \\
\le \;& \frac{d(q-1)}{2(i+2)}\{\sup_{v\in\mathbb{R}^{d-2}}|\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(v)|\}\Big\{E|E^{\mathcal{W}}(\tilde{S}_{i,1}\tilde{S}_{j,1})| + E|E^{\mathcal{W}}(\tilde{S}_{i,1}\tilde{S}_{j,2})| \\
& + E|E^{\mathcal{W}}(\tilde{S}_{i,1}S_{j,1})| + E|E^{\mathcal{W}}(\tilde{S}_{i,1}S_{j,2})| + E|E^{\mathcal{W}}(S_{i,1}\tilde{S}_{j,1})| \\
& + E|E^{\mathcal{W}}(S_{i,1}\tilde{S}_{j,2})| + E|E^{\mathcal{W}}(S_{i,1}S_{j,1})| + E|E^{\mathcal{W}}(S_{i,1}S_{j,2})|\Big\}.
\end{aligned}
$$

Now it follows from Lemma 1, Lemmas 7 to 10 and Lemma 12 that

$$
|\frac{d(q-1)}{4(i+2)}E[(\tilde{S}_i - S_i)(\tilde{S}_j - S_j)\frac{\partial^2}{\partial v_i \partial v_j}\psi_{\varepsilon^2}(V)]| \le O(\frac{\|h\|_\infty}{q^{1/2}})\log(1/\varepsilon),
$$

as $q \to \infty$ uniformly over $0 < \varepsilon < 1$. This proves Proposition 7. □

Lemma 12. *Let* $S_{\ell,1}$ *and* $\tilde{S}_{\ell,1}$, $\ell = 1, \dots, d-2$, *be as in (48). Then for* $1 \le \ell_1 \ne \ell_2 \le d-2$,

$$\begin{aligned}
\frac{d(q-1)}{2(\ell_1+2)} E|E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,1})| &= O(q^{-1/2}),\\
\frac{d(q-1)}{2(\ell_1+2)} E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,1})| &= O(q^{-1/2}), \qquad \text{as } q \to \infty.
\end{aligned}$$

Proof. We observe from (48) that for $1 \le \ell_1 < \ell_2 \le d-2$,

$$\begin{aligned}
& E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,1})\\
={}& E^{\mathcal{W}}\Big\{ \frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2} \sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
& \times \mathcal{I}\{J \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\mathcal{I}\{\pi_J(a_{i_1,J}) = B_1 = \pi_J(a_{i_2,J})\}\\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}\\
={}& \frac{1}{dq^5\sigma_{\ell_1}\sigma_{\ell_2}} \sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2} \sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
& \times \sum_{k=1}^{d} \mathcal{I}\{k \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\mathcal{I}\{a_{i_1,k} = a_{i_2,k}\}\\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\}.
\end{aligned}$$

Hence

$$\begin{aligned}
& E\{[E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,1})]^2\}\\
={}& E\Big\{ \frac{1}{d^2q^{10}\sigma_{\ell_1}^2\sigma_{\ell_2}^2} \sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2} \sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\\
& \times \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\}\\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\dots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}\\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}
\end{aligned}$$

$$
\begin{aligned}
= \quad & R^{\dagger}_{\{1,2,3,4\}} + R^{\dagger}_{\{1,2,3\},\{4\}} + R^{\dagger}_{\{1,2,4\},\{3\}} + R^{\dagger}_{\{1,3,4\},\{2\}} + R^{\dagger}_{\{2,3,4\},\{1\}} \\
& + R^{\dagger}_{\{1,2\},\{3\},\{4\}} + R^{\dagger}_{\{1,3\},\{2\},\{4\}} + R^{\dagger}_{\{1,4\},\{2\},\{3\}} + R^{\dagger}_{\{2,3\},\{1\},\{4\}} + R^{\dagger}_{\{2,4\},\{1\},\{3\}} \\
& + R^{\dagger}_{\{3,4\},\{1\},\{2\}} + R^{\dagger}_{\{1,2\},\{3,4\}} + R^{\dagger}_{\{1,3\},\{2,4\}} + R^{\dagger}_{\{1,4\},\{2,3\}} + R^{\dagger}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
& R^{\dagger}_{\{1,2,3,4\}} \\
= \quad & E\Big\{\frac{1}{d^2 q^{10}\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=i_2=i_3=i_4=1}^{q^2} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \\
& \times \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & O(\frac{1}{q^4}), \\
& R^{\dagger}_{\{1,2,3\},\{4\}} \;=\; R^{\dagger}_{\{1,2,4\},\{3\}} \;=\; R^{\dagger}_{\{1,3,4\},\{2\}} \;=\; R^{\dagger}_{\{2,3,4\},\{1\}} \\
= \quad & E\Big\{\frac{1}{d^2 q^{10}\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=i_2=i_3=1}^{q^2} \sum_{1\le i_4\le q^2:i_4\ne i_1} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \\
& \times \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\ldots,j_{2,|u^{(2)}|}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\mathcal{I}\{a_{i_1,k_3} = a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & O(\frac{1}{q^3}), \\
& R^{\dagger}_{\{1,2\},\{3\},\{4\}} \;=\; R^{\dagger}_{\{3,4\},\{1\},\{2\}}
\end{aligned}
$$

$$
\begin{aligned}
= \quad & E\Big\{ \frac{1}{d^2 q^{10} \sigma_{\ell_1}^2 \sigma_{\ell_2}^2} \sum_{i_1=i_2=1}^{q^2} \sum_{1 \le i_3 \le q^2 : i_3 \ne i_1} \sum_{1 \le i_4 \le q^2 : i_4 \ne i_1, i_3} \sum_{0 \le u_1^{(1)}, \dots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
& \times \sum_{0 \le u_1^{(2)}, \dots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \dots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \dots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \dots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \dots, j_{2,|u^{(2)}|}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \dots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \dots, j_{4,|u^{(4)}|}\} \mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \dots; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \dots; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \dots; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \dots; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})] \Big\} \\
= \quad & O(\frac{1}{q^3}), \\
& R^\dagger_{\{1,3\},\{2\},\{4\}} \; = \; R^\dagger_{\{1,4\},\{2\},\{3\}} \; = \; R^\dagger_{\{2,3\},\{1\},\{4\}} \; = \; R^\dagger_{\{2,4\},\{1\},\{3\}} \\
= \quad & E\Big\{ \frac{1}{d^2 q^{10} \sigma_{\ell_1}^2 \sigma_{\ell_2}^2} \sum_{i_1=i_3=1}^{q^2} \sum_{1 \le i_2 \le q^2 : i_2 \ne i_1} \sum_{1 \le i_4 \le q^2 : i_4 \ne i_1, i_2} \sum_{0 \le u_1^{(1)}, \dots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
& \times \sum_{0 \le u_1^{(2)}, \dots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \dots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \dots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \dots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \dots, j_{2,|u^{(2)}|}\} \mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \dots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \dots, j_{4,|u^{(4)}|}\} \mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \dots; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \dots; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \dots; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \dots; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})] \Big\} \\
= \quad & O(\frac{1}{q^4}), \\
& R^\dagger_{\{1,3\},\{2,4\}} \; = \; R^\dagger_{\{1,4\},\{2,3\}} \\
= \quad & E\Big\{ \frac{1}{d^2 q^{10} \sigma_{\ell_1}^2 \sigma_{\ell_2}^2} \sum_{i_1=i_3=1}^{q^2} \sum_{1 \le i_2 = i_4 \le q^2 : i_2 \ne i_1} \sum_{0 \le u_1^{(1)}, \dots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
& \times \sum_{0 \le u_1^{(2)}, \dots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \dots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \dots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2}
\end{aligned}
$$

$$
\begin{aligned}
&\times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \dots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \dots, j_{2,|u^{(2)}|}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\} \\
&\times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \dots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \dots, j_{4,|u^{(4)}|}\}\mathcal{I}\{a_{i_1,k_3} = a_{i_2,k_3}\} \\
&\times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \dots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \dots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \dots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \dots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & O(\frac{1}{q^3}), \\
& R^{\dagger}_{\{1\},\{2\},\{3\},\{4\}} \\
= \quad & E\Big\{\frac{1}{d^2 q^{10} \sigma^2_{\ell_1} \sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1 \le i_2 \le q^2 : i_2 \ne i_1} \sum_{1 \le i_3 \le q^2 : i_3 \ne i_1, i_2} \sum_{1 \le i_4 \le q^2 : i_4 \ne i_1, i_2, i_3} \sum_{0 \le u_1^{(1)}, \dots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
&\times \sum_{0 \le u_1^{(2)}, \dots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \dots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \dots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
&\times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \dots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \dots, j_{2,|u^{(2)}|}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\} \\
&\times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \dots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \dots, j_{4,|u^{(4)}|}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
&\times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \dots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}}); \dots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}}); \dots ; \pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}}); \dots ; \pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & O(\frac{1}{q^4}), \\
& R^{\dagger}_{\{1,2\},\{3,4\}} \\
= \quad & E\Big\{\frac{1}{d^2 q^{10} \sigma^2_{\ell_1} \sigma^2_{\ell_2}} \sum_{i_1=1}^{q^2} \sum_{1 \le i_3 \le q^2 : i_3 \ne i_1} \sum_{0 \le u_1^{(1)}, \dots, u_d^{(1)} \le 1 : |u^{(1)}| = \ell_1 + 2} \\
&\times \sum_{0 \le u_1^{(2)}, \dots, u_d^{(2)} \le 1 : |u^{(2)}| = \ell_2 + 2} \sum_{0 \le u_1^{(3)}, \dots, u_d^{(3)} \le 1 : |u^{(3)}| = \ell_1 + 2} \sum_{0 \le u_1^{(4)}, \dots, u_d^{(4)} \le 1 : |u^{(4)}| = \ell_2 + 2} \\
&\times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1}, \dots, j_{1,|u^{(1)}|}\} \cap \{j_{2,1}, \dots, j_{2,|u^{(2)}|}\} \\
&\times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1}, \dots, j_{3,|u^{(3)}|}\} \cap \{j_{4,1}, \dots, j_{4,|u^{(4)}|}\} \\
&\times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}}); \dots ; \pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\nu^*[\pi_{j_{2,1}}(a_{i_1,j_{2,1}}); \dots ; \pi_{j_{2,|u^{(2)}|}}(a_{i_1,j_{2,|u^{(2)}|}})]
\end{aligned}
$$

$$
\begin{aligned}
& \times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\nu^*[\pi_{j_{4,1}}(a_{i_3,j_{4,1}});\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_3,j_{4,|u^{(4)}|}})]\Big\} \\
= \quad & O(\frac{1}{q^3}),
\end{aligned}
$$

as $q \to \infty$. The last equality ues the assumption that $\ell_1 \neq \ell_2$. Thus we conclude that

$$
\frac{d(q-1)}{2(\ell_1+2)}E|E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,1})| \leq \frac{d(q-1)}{2(\ell_1+2)}\{E[E^{\mathcal{W}}(S_{\ell_1,1}S_{\ell_2,1})]^2\}^{1/2} = O(\frac{1}{q^{1/2}}),
$$

as $q \to \infty$. Next we observe that from (48) that for $1 \leq \ell_1 < \ell_2 \leq d-2$,

$$
\begin{aligned}
& E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,1}) \\
= \quad & E^{\mathcal{W}}\Big\{\frac{1}{q^4\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \\
& \times\mathcal{I}\{J \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{\pi_J(a_{i_1,J}) = B_1 = \pi_J(a_{i_2,J})\} \\
& \times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tau_{B_1,B_2}\circ\pi_J(a_{i_1,J});\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tau_{B_1,B_2}\circ\pi_J(a_{i_2,J});\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\Big\} \\
= \quad & \frac{1}{dq^5(q-1)\sigma_{\ell_1}\sigma_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2} \\
& \times\sum_{k=1}^{d}\mathcal{I}\{k \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k} = a_{i_2,k}\} \\
& \times\sum_{0\leq\tilde{c}_k\leq q-1:\tilde{c}_k\neq\pi_k(a_{i_1,k})}\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_k;\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_k;\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})],
\end{aligned}
$$

and

$$
\begin{aligned}
& E\{[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,1})]^2\} \\
= \quad & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=1}^{q^2}\sum_{i_2=1}^{q^2}\sum_{i_3=1}^{q^2}\sum_{i_4=1}^{q^2}\sum_{0\leq u_1^{(1)},\ldots,u_d^{(1)}\leq 1:|u^{(1)}|=\ell_1+2} \\
& \times\sum_{0\leq u_1^{(2)},\ldots,u_d^{(2)}\leq 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\leq u_1^{(3)},\ldots,u_d^{(3)}\leq 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\leq u_1^{(4)},\ldots,u_d^{(4)}\leq 1:|u^{(4)}|=\ell_2+2} \\
& \times\sum_{k_1=1}^{d}\mathcal{I}\{k_1 \in \{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1} = a_{i_2,k_1}\} \\
& \times\sum_{k_3=1}^{d}\mathcal{I}\{k_3 \in \{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\}
\end{aligned}
$$

$$
\begin{aligned}
& \times \sum_{0\le \tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq \pi_{k_1}(a_{i_1,k_1})} \sum_{0\le \tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq \pi_{k_3}(a_{i_3,k_3})} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
& \times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & R^{\ddagger}_{\{1,2,3,4\}} + R^{\ddagger}_{\{1,2,3\},\{4\}} + R^{\ddagger}_{\{1,2,4\},\{3\}} + R^{\ddagger}_{\{1,3,4\},\{2\}} + R^{\ddagger}_{\{2,3,4\},\{1\}} \\
& + R^{\ddagger}_{\{1,2\},\{3\},\{4\}} + R^{\ddagger}_{\{1,3\},\{2\},\{4\}} + R^{\ddagger}_{\{1,4\},\{2\},\{3\}} + R^{\dagger}_{\{2,3\},\{1\},\{4\}} + R^{\dagger}_{\{2,4\},\{1\},\{3\}} \\
& + R^{\ddagger}_{\{3,4\},\{1\},\{2\}} + R^{\ddagger}_{\{1,2\},\{3,4\}} + R^{\ddagger}_{\{1,3\},\{2,4\}} + R^{\ddagger}_{\{1,4\},\{2,3\}} + R^{\ddagger}_{\{1\},\{2\},\{3\},\{4\}},
\end{aligned}
$$

where

$$
\begin{aligned}
& R^{\ddagger}_{\{1,2,3,4\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i=i_2=i_3=i_4=1}^{q^2} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \\
& \times \sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2} \sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2} \sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
& \times \sum_{k_1=1}^{d} \mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\} \\
& \times \sum_{k_3=1}^{d} \mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\} \\
& \times \sum_{0\le \tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\neq \pi_{k_1}(a_{i_1,k_1})} \sum_{0\le \tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\neq \pi_{k_3}(a_{i_3,k_3})} \\
& \times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
& \times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
& \times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
& \times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \; & O(\frac{1}{q^4}), \\
& R^{\ddagger}_{\{1,2,3\},\{4\}} \; = \; R^{\ddagger}_{\{1,2,4\},\{3\}} \; = \; R^{\ddagger}_{\{1,3,4\},\{2\}} \; = \; R^{\ddagger}_{\{2,3,4\},\{1\}} \\
= \; & E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}} \sum_{i_1=i_2=i_3=1}^{q^2} \sum_{1\le i_4\le q^2:i_4\neq i_1} \sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}
\end{aligned}
$$

$$
\begin{aligned}
&\times \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
&\times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
&\times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\dots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
&\times \sum_{0\le \tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne \pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le \tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne \pi_{k_3}(a_{i_3,k_3})} \\
&\times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
&\times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
&\times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \ & O(\frac{1}{q^2}), \\
& R^{\ddagger}_{\{1,2\},\{3\},\{4\}} \ = \ R^{\ddagger}_{\{3,4\},\{1\},\{2\}} \\
= \ & E\Big\{\frac{1}{d^2 q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2} \sum_{i_1=i_2=1}^{q^2}\ \sum_{1\le i_3\le q^2:i_3\ne i_1}\ \sum_{1\le i_4\le q^2:i_4\ne i_1,i_3}\ \sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2} \\
&\times \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2} \\
&\times \sum_{k_1=1}^{d} \mathcal{I}\{k_1 \in \{j_{1,1},\dots,j_{1,|u^{(1)}|}\} \cap \{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\} \\
&\times \sum_{k_3=1}^{d} \mathcal{I}\{k_3 \in \{j_{3,1},\dots,j_{3,|u^{(3)}|}\} \cap \{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3} = a_{i_4,k_3}\} \\
&\times \sum_{0\le \tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne \pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le \tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne \pi_{k_3}(a_{i_3,k_3})} \\
&\times \nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})] \\
&\times \nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})] \\
&\times \nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})] \\
&\times \nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\} \\
= \ & O(\frac{1}{q^3}), \\
& R^{\ddagger}_{\{1,3\},\{2\},\{4\}} \ = \ R^{\ddagger}_{\{1,4\},\{2\},\{3\}} \ = \ R^{\dagger}_{\{2,3\},\{1\},\{4\}} \ = \ R^{\dagger}_{\{2,4\},\{1\},\{3\}}
\end{aligned}
$$

$$
\begin{aligned}
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=i_3=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\quad\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\quad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\quad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}\\
&\quad\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\quad\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
&= O(\frac{1}{q^4}),\\
&\quad R^{\ddagger}_{\{1,3\},\{2,4\}} \;=\; R^{\ddagger}_{\{1,4\},\{2,3\}}\\
&= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma_{\ell_1}^2\sigma_{\ell_2}^2}\sum_{i_1=i_3=1}^{q^2}\sum_{1\le i_2=i_4\le q^2:i_2\ne i_3}\sum_{0\le u_1^{(1)},\ldots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\\
&\quad\times\sum_{0\le u_1^{(2)},\ldots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\sum_{0\le u_1^{(3)},\ldots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\sum_{0\le u_1^{(4)},\ldots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}\\
&\quad\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\ldots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\ldots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}\\
&\quad\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\ldots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\ldots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_1,k_3}=a_{i_2,k_3}\}\\
&\quad\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}\\
&\quad\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\quad\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\quad\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\quad\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}
\end{aligned}
$$

$$= O(\frac{1}{q^3}),$$

$$R^{\ddagger}_{\{1\},\{2\},\{3\},\{4\}}$$

$$= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=1}^{q^2}\sum_{1\le i_2\le q^2:i_2\ne i_1}\sum_{1\le i_3\le q^2:i_3\ne i_1,i_2}\sum_{1\le i_4\le q^2:i_4\ne i_1,i_2,i_3}$$

$$\times\sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}\ \sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}$$

$$\times\sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}$$

$$\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}\mathcal{I}\{a_{i_1,k_1}=a_{i_2,k_1}\}$$

$$\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}\mathcal{I}\{a_{i_3,k_3}=a_{i_4,k_3}\}$$

$$\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}$$

$$\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]$$

$$\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\dots;\tilde{c}_{k_1};\dots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]$$

$$\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]$$

$$\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\dots;\tilde{c}_{k_3};\dots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}$$

$$= O(\frac{1}{q^4}),$$

$$R^{\ddagger}_{\{1,2\},\{3,4\}}$$

$$= E\Big\{\frac{1}{d^2q^{10}(q-1)^2\sigma^2_{\ell_1}\sigma^2_{\ell_2}}\sum_{i_1=i_2=1}^{q^2}\ \sum_{1\le i_3=i_4\le q^2:i_3\ne i_2}\ \sum_{0\le u_1^{(1)},\dots,u_d^{(1)}\le 1:|u^{(1)}|=\ell_1+2}$$

$$\times\sum_{0\le u_1^{(2)},\dots,u_d^{(2)}\le 1:|u^{(2)}|=\ell_2+2}\ \sum_{0\le u_1^{(3)},\dots,u_d^{(3)}\le 1:|u^{(3)}|=\ell_1+2}\ \sum_{0\le u_1^{(4)},\dots,u_d^{(4)}\le 1:|u^{(4)}|=\ell_2+2}$$

$$\times\sum_{k_1=1}^{d}\mathcal{I}\{k_1\in\{j_{1,1},\dots,j_{1,|u^{(1)}|}\}\cap\{j_{2,1},\dots,j_{2,|u^{(2)}|}\}\}$$

$$\times\sum_{k_3=1}^{d}\mathcal{I}\{k_3\in\{j_{3,1},\dots,j_{3,|u^{(3)}|}\}\cap\{j_{4,1},\dots,j_{4,|u^{(4)}|}\}\}$$

$$\times\sum_{0\le\tilde{c}_{k_1}\le q-1:\tilde{c}_{k_1}\ne\pi_{k_1}(a_{i_1,k_1})}\ \sum_{0\le\tilde{c}_{k_3}\le q-1:\tilde{c}_{k_3}\ne\pi_{k_3}(a_{i_3,k_3})}$$

$$
\begin{aligned}
&\times\nu^*[\pi_{j_{1,1}}(a_{i_1,j_{1,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{1,|u^{(1)}|}}(a_{i_1,j_{1,|u^{(1)}|}})]\\
&\times\nu^*[\pi_{j_{2,1}}(a_{i_2,j_{2,1}});\ldots;\tilde{c}_{k_1};\ldots;\pi_{j_{2,|u^{(2)}|}}(a_{i_2,j_{2,|u^{(2)}|}})]\\
&\times\nu^*[\pi_{j_{3,1}}(a_{i_3,j_{3,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{3,|u^{(3)}|}}(a_{i_3,j_{3,|u^{(3)}|}})]\\
&\times\nu^*[\pi_{j_{4,1}}(a_{i_4,j_{4,1}});\ldots;\tilde{c}_{k_3};\ldots;\pi_{j_{4,|u^{(4)}|}}(a_{i_4,j_{4,|u^{(4)}|}})]\Big\}\\
= \;& O(\frac{1}{q^3}),
\end{aligned}
$$

as $q \to \infty$. The last equality uses the assumption that $\ell_1 < \ell_2$. Thus we conclude that

$$\frac{d(q-1)}{2(\ell_1+2)}E|E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,1})| \leq \frac{d(q-1)}{2(\ell_1+2)}\{E[E^{\mathcal{W}}(\tilde{S}_{\ell_1,1}\tilde{S}_{\ell_2,1})]^2\}^{1/2} = O(\frac{1}{q^{1/2}}),$$

as $q \to \infty$. This proves Lemma 12. □

Proof of Theorem 2. Let Z_1 be a random variable having the standard (univariate) normal distribution and $\xi = (\sigma_1/\sigma, \ldots, \sigma_{d-2}/\sigma)'$. Then $\|\xi\|^2 = 1$ and $W = \xi' V$. For ease of exposition in the subsequent argument, we shall write $\xi = \xi(q)$ and $V = V(q)$.

We claim that $\xi' V \to Z_1$ in distribution as $q \to \infty$. We shall prove this claim by contraposition. Suppose the claim is false. Then there exists an interval, say $[a, b) \subset \mathbb{R}$, such that $P(\xi' V \in [a, b))$ does not converge to $P(Z_1 \in [a, b))$ as $q \to \infty$. Since $P(\xi' V \in [a, b)) \in [0, 1]$, by the compactness of $[0, 1]$, there exists a subsequence, say $\xi'(q_k)V(q_k)$, of $\xi' V$ such that $P(\xi'(q_k)V(q_k) \in [a, b))$ converges to a number, say $L \neq P(Z_1 \in [a, b))$. As $\|\xi(q_k)\| = 1$, there exists a further subsequence, say $\xi'(q_{k_l})V(q_{k_l})$, of $\xi'(q_k)V(q_k)$ such that $\xi(q_{k_l})$ converges to a point $\tilde{\xi} \in \mathbb{R}^{d-2}$ as $q_{k_l} \to \infty$. This implies that $\xi'(q_{k_l})V(q_{k_l}) - \tilde{\xi}' V(q_{k_l}) \to 0$ in probability as $q_{k_l} \to \infty$ and hence $\xi'(q_{k_l})V(q_{k_l})$ and $\tilde{\xi}' V(q_{k_l})$ have the same asymptotic distribution. Using Theorem 3 and $\|\tilde{\xi}\|^2 = 1$, we observe that $\tilde{\xi}' V(q_{k_l})$ converges in law to the standard normal distribution as $q_{k_l} \to \infty$. Hence $\xi'(q_{k_l})V(q_{k_l})$ converges in law to the same latter distribution. This is a contradiction and the claim is proved.

We observe from Theorem 1 and Proposition 1 that for $\int_{[0,1)^d} f^2_{rem}(x)dx > 0$, we have $\sigma^2_{oal}/\sigma^2 = 1 + O(q^{-1})$ and $\sigma^2_{oas}/\sigma^2 = 1 + O(q^{-1})$ as $q \to \infty$. Thus we conclude from Proposition 2 and Slutsky's theorem that W_{oal} and W_{oas} both tend in law to the standard (univariate) normal distribution as $q \to \infty$. Finally using Theorem 1 and Proposition 1, we have $\lim_{q\to\infty} \sigma^*_{oal}/\sigma_{oal} = 1$. Hence for $[a, b) \subset \mathbb{R}$,

$$P(W^*_{oal} \in [a, b)) = \frac{1}{d!}\sum\nolimits^* P(W_{oal} \in [a, b)) + o(1) \to P(Z_1 \in [a, b)),$$

as $q \to \infty$ where $\sum^*$ denotes summation over all the $d!$ permutations of the columns of A^{**}. This proves that W^*_{oal} converges in law to the standard normal distribution. The proof of Theorem 2 is complete. □

References

[1] Bhattacharya, R. N. and Rao, R. R. (1986). *Normal Approximations and Asymptotic Expansions.* Krieger, Malabar.

[2] Bolthausen, E. and Götze, F. (1993). The rate of convergence for multivariate sampling statistics. *Ann. Statist.* **21** 1692-1710.

[3] Davis, P. J. and Rabinowitz, P. (1984). *Methods of Numerical Integration*, 2nd edition. Academic Press, Orlando.

[4] Götze, F. (1991). On the rate of convergence in the multivariate CLT. *Ann. Probab.* **19** 724-739.

[5] Hedayat, A. S., Sloane, N. J. A. and Stufken, J. (1999). *Orthogonal Arrays: Theory and Applications.* Springer, New York.

[6] Loh, W. L. (1996). A combinatorial central limit theorem for randomized orthogonal array sampling designs. *Ann. Statist.* **24** 1209-1224.

[7] McKay, M. D., Conover, W. J. and Beckman, R. J. (1979). A comparison of three methods for selecting values of output variables in the analysis of output from a computer code. *Technometrics* **21** 239-245.

[8] Owen, A. B. (1992a). Orthogonal arrays for computer experiments, integration and visualization. *Statist. Sinica* **2** 439-452.

[9] Owen, A. B. (1992b). A central limit theorem for Latin hypercube sampling. *J. Roy. Statist. Soc. Ser. B* **54** 541-551.

[10] Owen, A. B. (1994). Lattice sampling revisited: Monte Carlo variance of means over randomized orthogonal arrays. *Ann. Statist.* **22** 930-945.

[11] Owen, A. B. (1995). Randomly permuted (t, m, s)-nets and (t, s)-sequences. *Monte Carlo and Quasi-Monte Carlo Methods in Scientific Computing. Lecture Notes in Statist.* **106** 299-317. Springer, New York.

[12] Owen, A. B. (1997a). Monte Carlo variance of scrambled net quadrature. *SIAM J. Numer. Anal.* **34** 1884-1910.

[13] Owen, A. B. (1997b). Scrambled net variance for integrals of smooth functions. *Ann. Statist.* **25** 1541-1562.

[14] Raghavarao, D. (1971). *Constructions and Combinatorial Problems in Design of Experiments.* Wiley, New York.

[15] Sacks, J., Welch, W. J., Mitchell, T. J. and Wynn, H. P. (1989). Design and analysis of computer experiments. *Statist. Sci.* **4** 409-423.

[16] Santner, T. J., Williams, B. J. and Notz, W. I. (2003). *The Design and Analysis of Computer Experiments.* Springer, New York.

[17] Stein, C. M. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. *Proc. Sixth Berkeley Symp. Math. Statist. Probab.* **2** 583-602. Univ. California Press, Berkeley.

[18] Stein, C. M. (1986). *Approximate Computation of Expectations. IMS Lecture Notes–Monograph Ser.* **7**. IMS, Hayward, California.

[19] Stein, M. L. (1987). Large sample properties of simulations using Latin hypercube sampling. *Technometrics* **29** 143-151.

[20] Tang, B. (1993). Orthogonal array-based Latin hypercubes. *J. Amer. Statist. Assoc.* **88** 1392-1397.